

# MAT 2114 Intro to Linear Algebra

Joe Wells  
Virginia Tech

Spring 2020

## Contents

|   |    |
|---|----|
| Preface . . . . .                                       | 4  |
| 1.1 The Geometry and Algebra of Vectors . . . . .       | 5  |
| 1.1.1 Vector operations . . . . .                       | 6  |
| 1.1.2 Geometry of vector operations . . . . .           | 6  |
| 1.1.3 Linear combinations . . . . .                     | 7  |
| Exercises . . . . .                                     | 10 |
| 3.1 Matrix Operations . . . . .                         | 12 |
| 3.1.1 Matrix Basics . . . . .                           | 12 |
| 3.1.2 Matrix Operations . . . . .                       | 13 |
| 3.1.3 Matrix Powers . . . . .                           | 15 |
| 3.1.4 Transpose . . . . .                               | 15 |
| Exercises . . . . .                                     | 17 |
| 3.2 Matrix Algebra . . . . .                            | 18 |
| 3.2.1 Properties of Matrix Multiplication . . . . .     | 18 |
| Exercises . . . . .                                     | 21 |
| 2.1 Introduction to Linear Systems . . . . .            | 22 |
| Exercises . . . . .                                     | 25 |
| 2.2 Direct Methods for Solving Linear Systems . . . . . | 26 |
| Exercises . . . . .                                     | 32 |
| 2.3 Spanning Sets and Linear Independence . . . . .     | 34 |
| Exercises . . . . .                                     | 39 |

|                                       |   |    |
|---------------------------------------|---|----|
| 3.3                                   | The Inverse of a Matrix . . . . .                               | 40 |
|                                       | Exercises . . . . .   | 45 |
| 3.4                                   | Subspaces, Basis, Dimension, and Rank . . . . .                 | 46 |
| 3.4.1                                 | Subspaces Associated with Matrices . . . . .                    | 47 |
| 3.4.2                                 | Dimension and Rank . . . . .                                    | 51 |
| 3.4.3                                 | Coordinates . . . . .   | 53 |
|                                       | Exercises . . . . .   | 55 |
| 6.3                                   | Change of Basis . . . . .                                       | 56 |
| 3.6                                   | Introduction to Linear Transformations . . . . .                | 57 |
| 3.6.1                                 | Types of Linear Transformations of $\mathbb{R}^2$ . . . . .     | 58 |
| 3.7                                   | Applications . . . . .  | 61 |
| 3.7.1                                 | Markov Chains . . . . .   | 61 |
| 4.1                                   | Introduction to Eigenvalues and Eigenvectors . . . . .          | 63 |
| Appendix C: Complex Numbers . . . . . |   | 66 |
|                                       | Matrices with complex entries . . . . .                         | 66 |
| 4.2                                   | Determinants . . . . .  | 67 |
| 4.2.1                                 | Determinant of a $2 \times 2$ Matrix . . . . .                  | 67 |
| 4.2.2                                 | Determinant of a $n \times n$ Matrix . . . . .                  | 69 |
| 4.2.3                                 | Properties of Determinants . . . . .                            | 70 |
| 4.3                                   | Eigenvalues and Eigenvectors of $n \times n$ Matrices . . . . . | 72 |
| 4.3.1                                 | Relationship to Matrix Operations . . . . .                     | 73 |
| 4.4                                   | Similarity and Diagonalization . . . . .                        | 76 |
| 4.4.1                                 | Properties of similarity and similar matrices . . . . .         | 76 |
| 4.4.2                                 | Diagonalization . . . . .                                       | 77 |
| 4.4.3                                 | Computational power of diagonal matrices . . . . .              | 80 |
| 1.2                                   | Length and Angle: The Dot Product . . . . .                     | 81 |
| 1.2.1                                 | Length . . . . .  | 81 |
| 1.2.2                                 | Distances . . . . .   | 83 |
| 1.2.3                                 | Projections . . . . .   | 84 |
| 5.1                                   | Orthogonality in $\mathbb{R}^n$ . . . . .                       | 86 |
| 5.2                                   | Orthogonal Complements and Orthogonal Projections . . . . .     | 87 |

|     |   |    |
|-----|---|----|
| 5.3 | The Gram-Schmidt Process and the $QR$ Factorization . . . . . | 88 |
| 7.3 | Least Squares . . . . .                                       | 89 |

## Preface


There are many different approaches to linear algebra, and everyone has their preference. This document is compiled from the course I taught in the Spring of 2020 at Virginia Tech, where both the book (*Linear Algebra: A Modern Introduction* 4th Ed. by David Poole) and order of topics covers were suggested to me by some others in the department. Although not formally stated anywhere, this class was largely geared towards math-adjacent students (engineering, physics, computer science, etc.) and so these notes and the presentation are at a lower level of abstraction (and occasionally rigor) than what one might experience in another introductory linear algebra course. In hindsight, I probably would have picked both a different text and order in which to introduce the topics – it seems perverse to leave the phrase “vector space” until the 6<sup>th</sup> chapter! Nevertheless, I did my best to gently introduce concepts as needed in order to more smoothly segue the topics, and many of the homework exercises are designed to bridge certain theoretical gaps in the material (which is precisely where I’ve hidden the definition of a vector space).

I would like to thank the many students who inadvertently served as my copy editors for the semester.

## 1.1 The Geometry and Algebra of Vectors

$A = (a_1, \dots, a_n)$  and  $B = (b_1, \dots, b_n)$  points in space.

**Definition.** The *vector*  $\mathbf{AB}$  is the arrow from  $A$  (the *initial point* or *tail*) to  $B$  (the *terminal point* or *head*). In coordinates, use square brackets and write  $\mathbf{AB} = [b_1 - a_1, \dots, b_n - a_n]$ , and the individual coordinates of this vector are called *components*. A vector  $\mathbf{AB}$  is in *standard position* if  $A$  is at the origin. The *zero vector*, denoted  $\mathbf{0}$ , is the vector of all zeros.



*Remark.* Unless otherwise specified, given an arbitrary vector  $\mathbf{v}$ , we will always draw it in standard position.

This leads us to the following fact:

*Fact.* Two vectors are *equal* if and only if their components are the same.

We will use the following notation repeatedly throughout the semester:

- $\mathbb{R}$  - This is the set of real numbers.
- $\mathbb{R}^n$  - This is the set of vectors with  $n$  real components.

There are different ways to represent the vector  $\mathbf{v}$ , either as a *row vector*,

$$\mathbf{v} = [v_1, \dots, v_n],$$

or as a *column vector*,

$$\mathbf{v} = \begin{bmatrix} v_1 \\ \vdots \\ v_n \end{bmatrix}.$$

As will be come apparent in future sections, we'll eventually come to prefer column vectors, but for now we won't make a distinction.

*Remark.* Since writing a column vector takes up a lot of vertical space on the page, we will sometimes write  $\mathbf{v} = [v_1, \dots, v_n]^T$  to denote the column form of the vector  $\mathbf{v}$  (and we will also formally describe the superscript  $T$  notation in a later section).

### 1.1.1 Vector operations

Because the real numbers have proven to be so nice to us over the years, we would like to try to come up with an analogous algebraic structure that works for higher dimensions. The structure we're building is formally known as a *(real) vector space*, but we'll avoid that terminology and generality in favor of trying to motivate concrete understanding.

**Definition.** There are two operations on  $\mathbb{R}^n$ :

- *Vector addition* – For vectors  $\mathbf{u} = [u_1, \dots, u_n]$  and  $\mathbf{v} = [v_1, \dots, v_n]$  in  $\mathbb{R}^n$

$$\mathbf{u} + \mathbf{v} = [u_1 + v_1, \dots, u_n + v_n]$$

- *Scalar multiplication.* For a vector  $\mathbf{u} = [u_1, \dots, u_n]$  in  $\mathbb{R}^n$  and a *scalar*  $a$  in  $\mathbb{R}$ ,

$$a\mathbf{u} = [au_1, \dots, au_n]$$

It's worthwhile to note that scalar multiplication is slightly weaker than what we would want – ideally we could multiply two vectors together to get a vector, but instead we can only multiply a vector and a scalar (at least the output is still a vector). The reasons for this are somewhat subtle and deep, but I hope in a later section to provide some motivation for why trying to define other types of multiplication is ultimately more problematic.

**Theorem 1.1.1** (*Poole Theorem 1.1 - Algebraic Properties of Vectors in  $\mathbb{R}^n$* ). *Let  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  be vectors in  $\mathbb{R}^n$  and let  $a, b$  be scalars in  $\mathbb{R}$ . With vector addition and scalar multiplication on  $\mathbb{R}^n$  as defined above, the following are true:*

- $\mathbf{u} + \mathbf{v}$  is in  $\mathbb{R}^n$ .
- $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$ .
- $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$ .
- The zero vector  $\mathbf{0}$  satisfies  $\mathbf{u} + \mathbf{0} = \mathbf{u}$  for all vectors  $\mathbf{u}$ .
- For each  $\mathbf{u}$  in  $\mathbb{R}^n$ , there is some vector  $-\mathbf{u}$  for which  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .
- $a\mathbf{u}$  is in  $\mathbb{R}^n$ .
- $a(\mathbf{u} + \mathbf{v}) = a\mathbf{u} + b\mathbf{v}$ .
- $(a + b)\mathbf{u} = a\mathbf{u} + b\mathbf{u}$ .
- $(ab)\mathbf{u} = a(b\mathbf{u})$ .
- $1\mathbf{u} = \mathbf{u}$ .

All parts of the above theorem are straightforward to check, so they are left as an exercise to the reader. The above theorem effectively says that  $\mathbb{R}^n$ , with our two operations, satisfies all of the same algebraic properties as  $\mathbb{R}$ .

### 1.1.2 Geometry of vector operations

Given two vectors  $\mathbf{u}$  and  $\mathbf{v}$ , their sum is the vector formed from adjoining the tail of  $\mathbf{v}$  to the head of  $\mathbf{u}$ . In  $\mathbb{R}^2$ , this is equivalent to the diagonal to the parallelogram formed from sides  $\mathbf{u}$  and  $\mathbf{v}$ . Scalar multiplication merely *scales* the vector and leaves its direction unchanged. To see this, consider the following example:


**Example 1.1.2.** Let  $\mathbf{u} = [-1, -2]$ ,  $\mathbf{v} = [4, 1]$ , and  $a = 2$ . Then

$$\mathbf{u} + \mathbf{v} = [-1 + 4, -2 + 1] = [3, -1]$$

and

$$a\mathbf{u} = [2(-1), 2(-2)] = [-2, -4].$$

Drawing them out



### 1.1.3 Linear combinations

**Definition.** A vector  $\mathbf{u}$  is a *linear combination* of the vectors  $\mathbf{v}_1, \dots, \mathbf{v}_n$  if there are scalars  $a_1, \dots, a_n$  so that

$$\mathbf{u} = a_1\mathbf{v}_1 + \dots + a_n\mathbf{v}_n.$$

You can think of a linear combination as some sort of recipe - the  $\mathbf{v}_i$ 's are the ingredients, the  $a_i$ 's are the quantities of those ingredients, and  $\mathbf{u}$  is the finished product.

**Definition.** In  $\mathbb{R}^n$ , there are  $n$  vectors

$$\begin{aligned} \mathbf{e}_1 &= [1, 0, \dots, 0], \\ \mathbf{e}_2 &= [0, 1, \dots, 0], \\ &\vdots \\ \mathbf{e}_n &= [0, 0, \dots, 1], \end{aligned}$$

which we call the *standard basis vectors* for  $\mathbb{R}^n$ .

For now, ignore the word *basis* above; we will give technical meaning to that later. The reason these are standard is because, when looking to decompose a vector  $\mathbf{u}$  into a linear combination of vectors, then simply picking apart the components is probably the most natural thing to try first.


**Example 1.1.3.** The vector  $\mathbf{u} = \begin{bmatrix} 5 \\ 6 \\ 7 \end{bmatrix}$  is a linear combination of the standard basis vectors in the following way:

$$\mathbf{u} = \begin{bmatrix} 5 \\ 6 \\ 7 \end{bmatrix} = \begin{bmatrix} 5 \\ 0 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 6 \\ 0 \end{bmatrix} + \begin{bmatrix} 0 \\ 0 \\ 7 \end{bmatrix} = 5 \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} + 6 \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} + 7 \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} = 5\mathbf{e}_1 + 6\mathbf{e}_2 + 7\mathbf{e}_3$$


**Definition.** A *coordinate grid* is the grid formed by translates of a set of vectors.

I don't think the above definition is a very good way to explain the idea, which is actually fairly intuitive. We want to form a grid where (1) the lines parallel to the given vectors and (2) the intersections of these grids correspond to integer-linear combinations of vectors. It may be more enlightening to just see a few examples.

**Example 1.1.4.** The coordinate grid for  $\mathbb{R}^2$  formed from the standard basis vectors  $\mathbf{e}_1$  and  $\mathbf{e}_2$  is the usual Cartesian grid.



**Example 1.1.5.** The coordinate grid for  $\mathbb{R}^2$  formed from the vectors  $\mathbf{v}_1 = [1, 1]$  and  $\mathbf{v}_2 = [1, -1]$  is below.





Combined with the geometric intuition about vector addition and scalar multiplication, these coordinate grids provide us with a way to visually identify the linear combination.



**Example 1.1.6.** Write the vector  $\mathbf{u} = [2, 4]$  as a linear combination of the vectors...

1.  $\mathbf{e}_1 = [1, 0]$ ,  $\mathbf{e}_2 = [0, 1]$
2.  $\mathbf{v}_1 = [1, 1]$ ,  $\mathbf{v}_2 = [1, -1]$



We can verify this algebraically:

$$\begin{aligned}\mathbf{u} &= 2\mathbf{e}_1 + 4\mathbf{e}_2 \\ &= 2[1, 0] + 4[0, 1] \\ &= [2, 0] + [0, 4] \\ &= [2, 4]\end{aligned}$$

We can verify this algebraically:

$$\begin{aligned}\mathbf{u} &= 3\mathbf{v}_1 + (-1)\mathbf{v}_2 \\ &= 3[1, 1] + (-1)[1, -1] \\ &= [3, 3] + [-1, 1] \\ &= [2, 4]\end{aligned}$$

## Section 1.1 Exercises

1. Draw and label the following vectors in standard position.

$$(a) \mathbf{u} = \begin{bmatrix} 4 \\ -1 \end{bmatrix}$$

$$(b) \mathbf{v} = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$$

$$(c) \mathbf{w} = \begin{bmatrix} -3 \\ 2 \end{bmatrix}$$

2. Solve for the vector  $\mathbf{x}$  satisfying the following equation:  $2\mathbf{x} + \begin{bmatrix} 5 \\ 7 \\ 1 \end{bmatrix} = 3 \left( \mathbf{x} + \begin{bmatrix} -3 \\ -1 \\ 0 \end{bmatrix} \right)$

3. Sketch the coordinate grid formed by the vectors  $\mathbf{u} = [-1, 2]$ ,  $\mathbf{v} = [-2, -1]$  and use it to draw the vector  $\mathbf{w} = \mathbf{u} - 2\mathbf{v}$ .

4. (a) Sketch the coordinate grid formed by the vectors  $\mathbf{u} = \begin{bmatrix} 1 \\ -3 \end{bmatrix}$  and  $\mathbf{v} = \begin{bmatrix} -1 \\ 1 \end{bmatrix}$ . Use this to

write  $\mathbf{x} = \begin{bmatrix} 2 \\ -5 \end{bmatrix}$  as a linear combination of  $\mathbf{u}$  and  $\mathbf{v}$ . Verify your answer algebraically.

(b) Write  $\mathbf{x} = \begin{bmatrix} 2 \\ -5 \\ 0 \end{bmatrix}$  as a linear combination of  $\mathbf{u} = \begin{bmatrix} 1 \\ -3 \\ 0 \end{bmatrix}$ ,  $\mathbf{v} = \begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ , and  $\mathbf{w} = \begin{bmatrix} 3 \\ 1 \\ 1 \end{bmatrix}$ .

**Definition.** A (real) vector space  $V$  is a collection of objects (called *vectors*) with two operations *vector addition* (denoted  $+$ ) and *scalar multiplication* (no symbol) satisfying the following properties: For all vectors  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  and real numbers  $a, b$  (called *scalars*),

(a)  $\mathbf{u} + \mathbf{v}$  is in  $V$  [closure of addition]

(b)  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$  [commutativity of addition]

(c)  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  [associativity of  $+$ ]

(d) There is some vector  $\mathbf{0}$ , called the *zero vector*, so that  $\mathbf{u} + \mathbf{0} = \mathbf{u}$  for all vectors  $\mathbf{u}$ . [additive identity]

(e) For each  $\mathbf{u}$  in  $V$ , there is some vector  $-\mathbf{u}$  for which  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ . [additive inverse]

(f)  $a\mathbf{u}$  is in  $V$  [closure of scalar mult.]

(g)  $a(\mathbf{u} + \mathbf{v}) = a\mathbf{u} + a\mathbf{v}$  [distributivity]

(h)  $(a + b)\mathbf{u} = a\mathbf{u} + b\mathbf{u}$  [distributivity]

(i)  $(ab)\mathbf{u} = a(b\mathbf{u})$  [associativity of scalar mult.]

(j)  $1\mathbf{u} = \mathbf{u}$  [scalar mult. identity]

For example,  $\mathbb{R}^n$  is a real vector space (this is just Theorem 1.1.1).

5. **Challenge.** Let  $\mathcal{P}^2$  be the collection of polynomials with indeterminate  $x$ , with real numbered coefficients, and with degree 2 (or less). Let

$$\begin{aligned} \mathbf{u} &= u_0 + u_1x + u_2x^2, \\ \mathbf{v} &= v_0 + v_1x + v_2x^2, \quad \text{and} \\ \mathbf{w} &= w_0 + w_1x + w_2x^2 \end{aligned}$$

be arbitrary polynomials in  $\mathcal{P}^2$  and let  $a, b$  be scalars in  $\mathbb{R}$ . Define the following operations on  $\mathcal{P}^2$ :

- vector addition:

$$\mathbf{u} + \mathbf{v} = (u_0 + v_0) + (u_1 + v_1)x + (u_2 + v_2)x^2$$

- scalar multiplication:

$$a\mathbf{u} = au_0 + au_1x + au_2x^2$$

Show that  $\mathcal{P}^2$  is a real vector space, i.e., show it satisfies axioms (a) - (j) above.

## 3.1 Matrix Operations

### 3.1.1 Matrix Basics

**Definition.** A *matrix* is an array of numbers (called *entries*) and has *size*  $m \times n$  if it has  $m$  rows and  $n$  columns.

$$A = [a_{ij}] = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \vdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix}$$

The subscripts on the entries  $a_{ij}$  tell us that we're looking at the entry in the  $i^{\text{th}}$  row and the  $j^{\text{th}}$  column.

*Fact.* Two matrices are equal if and only if both (1) their sizes are equal and (2) their corresponding entries are all equal.

**Example 3.1.1.** Consider the matrices

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 2 \\ 4 & 5 \end{bmatrix}, \quad \text{and } C = \begin{bmatrix} 7 & 8 \\ 9 & 10 \end{bmatrix}.$$

$A \neq B$  because their sizes are different, and  $A \neq C$  because their corresponding entries are not equal.

**Definition.** A matrix is *square* if it has size  $n \times n$  and the *diagonal of the matrix*  $A = [a_{ij}]$  are the entries where  $i = j$ .

$$A = \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & \ddots & \vdots \\ a_{n1} & \cdots & a_{nn} \end{bmatrix}$$

A square matrix is *diagonal* if the only nonzero entries are along the diagonal. You may see this written as  $A = \text{diag}(a_{11}, \dots, a_{nn})$ .

A bit of a subtlety – the definition of diagonal just says that nonzero entries must occur along the diagonal, but the diagonal entries *do not necessarily need to be nonzero*.

**Example 3.1.2.** The matrices

$$A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 7 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

are both diagonal because every entry off of the diagonal is 0.

**Definition.** A matrix is *scalar* if it is a diagonal matrix and the diagonal entries are all equal, i.e., the matrix  $A = \text{diag}(r, r, \dots, r)$  for some  $r \in \mathbb{R}$ .

**Definition.** The *zero matrix*, often denoted  $O$  is the matrix for which all entries are 0. Its size should be clear from context, but we may write  $O_{m \times n}$  if we need to specify.

**Definition.** The *Kroenecker delta*, denoted  $\delta_{ij}$ ,  $\delta_i^j$ , or  $\delta^{ij}$ , is the following:

$$\delta_{ij} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

**Definition.** The *identity matrix*  $I_n$  is the diagonal  $n \times n$  matrix with all 1's along the diagonal. You may sometimes see this written as  $I_n = [\delta_{ij}]$ .

**Example 3.1.3.** It may be useful to see exactly how the Kroenecker delta leads to the identity matrix.

$$I_3 = [\delta_{ij}] = \begin{bmatrix} \delta_{11} & \delta_{12} & \delta_{13} \\ \delta_{21} & \delta_{22} & \delta_{23} \\ \delta_{31} & \delta_{32} & \delta_{33} \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

### 3.1.2 Matrix Operations

**Definition.** Given two  $m \times n$  matrices  $A = [a_{ij}]$  and  $B = [b_{ij}]$ , the *matrix sum* is

$$A + B = [a_{ij} + b_{ij}] = \begin{bmatrix} a_{11} + b_{11} & \cdots & a_{1n} + b_{1n} \\ \vdots & \ddots & \vdots \\ a_{m1} + b_{m1} & \cdots & a_{mn} + b_{mn} \end{bmatrix}$$

**Example 3.1.4.** For  $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$  and  $B = \begin{bmatrix} 7 & 8 & 9 \\ 10 & 11 & 12 \end{bmatrix}$ ,

$$A + B = \begin{bmatrix} 1+7 & 2+8 & 3+9 \\ 4+10 & 5+11 & 6+12 \end{bmatrix} = \begin{bmatrix} 8 & 10 & 12 \\ 14 & 16 & 18 \end{bmatrix}.$$

*Remark.* When adding two matrices, they must have the same size.

**Definition.** For a matrix  $A = [a_{ij}]$  and a scalar  $r$ , the *scalar multiple of A* is the matrix

$$rA = [ra_{ij}]$$

**Example 3.1.5.** Given the matrix  $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$  and the scalar  $r = 5$ ,

$$rA = \begin{bmatrix} 5(1) & 5(2) \\ 5(3) & 5(4) \end{bmatrix} = \begin{bmatrix} 5 & 10 \\ 15 & 20 \end{bmatrix}$$

**Definition.** Subtraction of matrices is then defined in the obvious way:  $A - B = A + (-1)B$ .

*Remark.* The zero matrix satisfies the properties you want it to:  $A + O = A$  and  $A - A = O$

**Definition.** Given two vectors in  $\mathbb{R}^n$ ,  $\mathbf{u} = [u_1, \dots, u_n]$  and  $\mathbf{v} = [v_1, \dots, v_n]$ , the *dot product of u and v* is the real number

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + u_2v_2 + \cdots + u_nv_n$$

**Proposition 3.1.6.** For vectors  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^n$  and a scalar  $k \in \mathbb{R}$ , the dot product has the following properties:

- (a)  $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
- (b)  $\mathbf{u} \cdot (\mathbf{v} + \mathbf{w}) = \mathbf{u} \cdot \mathbf{v} + \mathbf{u} \cdot \mathbf{w}$
- (c)  $k(\mathbf{u} \cdot \mathbf{v}) = (k\mathbf{u}) \cdot \mathbf{v} = \mathbf{u} \cdot (k\mathbf{v})$

**Exercise 3.1.1.** Prove Proposition 3.1.6.

**Definition.** Given an  $m \times n$  matrix  $A = [a_{ij}]$  and an  $n \times p$  matrix  $B = [b_{ij}]$ , the *matrix product* of  $A$  and  $B$  is

$$AB = [c_{ij}]$$

where

$$c_{ij} = a_{i1}b_{1j} + a_{i2}b_{2j} + \cdots + a_{in}b_{nj}$$

in other words, the  $(i, j)^{\text{th}}$  entry of  $AB$  is the dot product of the  $i^{\text{th}}$  row of  $A$  (thought of as a row vector) and the  $j^{\text{th}}$  of column  $B$  (thought of as a column vector).

**Example 3.1.7.** Let  $A$  and  $B$  be the following matrices

$$A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} = \begin{bmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \\ \mathbf{a}_3 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 7 & 8 \\ 9 & 10 \end{bmatrix} = \begin{bmatrix} \mathbf{b}_1 & \mathbf{b}_2 \end{bmatrix}$$

where the  $\mathbf{a}_i$ 's are vectors forming the rows in  $A$  and the  $\mathbf{b}_i$ 's are the vectors forming the columns of  $B$ . We have that

$$\begin{aligned} c_{11} &= \mathbf{a}_1 \cdot \mathbf{b}_1 = 1(7) + 2(9) = 23, & c_{12} &= \mathbf{a}_1 \cdot \mathbf{b}_2 = 1(8) + 2(10) = 28, \\ c_{21} &= \mathbf{a}_2 \cdot \mathbf{b}_1 = 3(7) + 4(9) = 57, & c_{22} &= \mathbf{a}_2 \cdot \mathbf{b}_2 = 3(8) + 4(10) = 64, \\ c_{31} &= \mathbf{a}_3 \cdot \mathbf{b}_1 = 5(7) + 6(9) = 89, & c_{32} &= \mathbf{a}_3 \cdot \mathbf{b}_2 = 5(8) + 6(10) = 100, \end{aligned}$$

and so the product matrix is

$$AB = [c_{ij}] = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \\ c_{31} & c_{32} \end{bmatrix} = \begin{bmatrix} 23 & 28 \\ 57 & 64 \\ 89 & 100 \end{bmatrix}.$$

Thinking of  $\mathbf{b}_1$  and  $\mathbf{b}_2$  as matrices of size  $3 \times 1$ , we may notice that the first column of  $AB$  is precisely  $A\mathbf{b}_1$  and the second column is  $A\mathbf{b}_2$ :

$$\begin{aligned} \begin{bmatrix} c_{11} \\ c_{21} \\ c_{31} \end{bmatrix} &= \begin{bmatrix} 1(7) + 2(9) \\ 3(7) + 4(9) \\ 5(7) + 6(9) \end{bmatrix} = 7 \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} + 9 \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix} = A\mathbf{b}_1. \\ \begin{bmatrix} c_{12} \\ c_{22} \\ c_{32} \end{bmatrix} &= \begin{bmatrix} 1(8) + 2(10) \\ 3(8) + 4(10) \\ 5(8) + 6(10) \end{bmatrix} = 8 \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} + 10 \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix} = A\mathbf{b}_2. \end{aligned}$$

As such, we have

$$AB = \begin{bmatrix} c_{11} & c_{12} \\ c_{21} & c_{22} \\ c_{31} & c_{32} \end{bmatrix} = \begin{bmatrix} A\mathbf{b}_1 & A\mathbf{b}_2 \end{bmatrix}$$

The matrix on the right is sometimes called the *matrix-column representation* of the product  $AB$ .

*Remark.* Two matrices  $A, B$  can be multiplied even if their sizes are different. As long as the number of columns of  $A$  is equal to the number of rows of  $B$ , then the product  $AB$  exists. Moreover

$$\begin{matrix} A & B & = & AB \\ m \times n & n \times p & & m \times p \end{matrix}$$

*Fact.* If  $A$  is an  $m \times n$  matrix, then we have that

$$I_m A = A \quad \text{and} \quad A I_n = A$$

and it is for this reason that we call  $I_n$  the identity matrix.

For now you'll have to take it on faith that our definition matrix multiplication, although odd and tedious, is actually the correct notion. The parallel story of matrices is that they correspond to certain functions on vector spaces, and matrix multiplication corresponds to function composition (which, as you can imagine, can be quite confusing). We'll fully make this connection later this semester.

### 3.1.3 Matrix Powers

If  $A$  is a square matrix, then for positive integers  $k$ , then we can define the power of a matrix in the intuitive way,

$$A^k = \underbrace{AA \cdots A}_{k \text{ factors}}$$

and the usual rules for exponents hold:

- $A^r A^s = A^{r+s}$
- $(A^r)^s = A^{rs}$

**Example 3.1.8.** If  $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ , then

$$A^2 = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 7 & 10 \\ 15 & 22 \end{bmatrix}$$

and

$$A^3 = A^2 A = \begin{bmatrix} 7 & 10 \\ 15 & 22 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} = \begin{bmatrix} 37 & 54 \\ 81 & 118 \end{bmatrix}$$

### 3.1.4 Transpose

**Definition.** If  $A = [a_{ij}]$  is an  $m \times n$  matrix, then its *transpose*,  $A^T = [a_{ji}]$  is the  $n \times m$  matrix formed by swapping the  $(i, j)^{\text{th}}$  and  $(j, i)^{\text{th}}$  entries of  $A$ .

Visually, the transpose amounts to flipping the matrix across the red line below

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \xrightarrow{\text{flip}} \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix} = A^T$$

**Definition.** A matrix  $A$  is *symmetric* if  $A = A^T$ .

**Example 3.1.9.**  $A = \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}$  is symmetric, while  $B = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$  is not.

**Example 3.1.10.** Let  $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix}$ . Then

$$\begin{aligned} A^T A &= \begin{bmatrix} 1 & 3 & 5 \\ 2 & 4 & 6 \end{bmatrix} \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix} \\ &= \begin{bmatrix} 1^2 + 3^2 + 5^2 & 1(2) + 3(4) + 5(6) \\ 2(1) + 4(3) + 6(5) & 2^2 + 4^2 + 6^2 \end{bmatrix} \\ &= \begin{bmatrix} 35 & 44 \\ 44 & 56 \end{bmatrix} \end{aligned}$$

What's interesting to notice is that, while  $A$  is not symmetric (and not even square),  $A^T A$  is symmetric (and hence also a square matrix).



## Section 3.1 Exercises

1. Let  $A, \dots, F$  be matrices with the given sizes

| matrix | $A$          | $B$          | $C$          | $D$          | $E$          | $F$          |
|--------|--------------|--------------|--------------|--------------|--------------|--------------|
| size   | $2 \times 2$ | $2 \times 3$ | $3 \times 2$ | $2 \times 2$ | $1 \times 2$ | $2 \times 1$ |

Determine the sizes of each of the following matrices, if possible. If it is not possible, explain why.

- $3D - 2A$
  - $D + BC$
  - $BB^T$
  - $B^T C^T - (CB)^T$
  - $DA - AD$
  - $(I_2 - D)^2$
2. Compute  $(A - I_2)^2$  where  $A = \begin{bmatrix} 1 & -7 \\ 0 & 1 \end{bmatrix}$ .
3. Give an example of a nonzero  $2 \times 2$  matrix  $A$  for which  $A^2 = O_{2 \times 2}$ . *Show some work to verify it has this property.*
4. Give an example of a nonzero  $3 \times 3$  matrix  $A$  for which  $A^3 = O_{3 \times 3}$ . *Show some work to verify it has this property.*
5. Let  $A = \begin{bmatrix} 0 & 2 \\ 0 & 0 \end{bmatrix}$ . Give an example of two  $2 \times 2$  matrices  $B$  and  $C$  such that  $AB = AC$  but  $B \neq C$ . *Show some work to verify it has this property.*
6. Give an example of two  $2 \times 2$  matrices  $A$  and  $B$  for which  $AB \neq BA$ . *Show some work to verify it has this property.*
7. Let  $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$ .
- Compute  $A^2$ ,  $A^3$ , and  $A^4$ .
  - What is  $A^n$  for some positive integer  $n$ ?

## 3.2 Matrix Algebra

We'll introduce another piece of notation (that I probably should have used in the previous section):

$\mathbb{R}^{m \times n}$  represents the collection of  $m \times n$  matrices with real number entries.

**Theorem 3.2.1** (*Poole Theorem 3.2 - Algebraic Properties of Matrix Addition and Scalar Multiplication*). *Let  $A, B, C \in \mathbb{R}^{m \times n}$  and let  $c, d \in \mathbb{R}$ . The following are true:*

(a)  $A + B = B + A$

(b)  $(A + B) + C = A + (B + C)$

(c)  $A + O_{m \times n} = A$

(d)  $A + (-A) = O_{m \times n}$

(e)  $c(A + B) = cA + cB$

(f)  $(c + d)A = cA + dA$

(g)  $c(dA) = (cd)A$

(h)  $1A = A$

*Remark.* In short, Theorem 3.2.1 above says that  $\mathbb{R}^{m \times n}$  is a real vector space (see page 10).

### 3.2.1 Properties of Matrix Multiplication

Matrix multiplication is *not commutative in general*, and it is often the case that  $AB \neq BA$ . This fact is clear if  $A \in \mathbb{R}^{m \times n}$  and  $B \in \mathbb{R}^{n \times m}$  where  $m \neq n$  (just compare the sizes of  $AB$  and  $BA$ ), but is possibly less obvious in the case where  $A, B$  are both square matrices. It is an exercise to find an example of this in the case of  $2 \times 2$  matrices.

So what properties does matrix multiplication have?

**Theorem 3.2.2** (*Poole Theorem 3.3 - Properties of Matrix Multiplication*). *Let  $A, B, C$  be matrices (whose sizes are such that the following exist) and  $k \in \mathbb{R}$  a scalar. Then*

(a)  $A(BC) = (AB)C$

(b)  $A(B + C) = AB + AC$

(c)  $(A + B)C = AC + BC$

(d)  $k(AB) = (kA)B = A(kB)$

(e)  $I_m A = A = A I_n$  (if  $A$  is  $m \times n$ )

*Remark.* This theorem implies that  $\mathbb{R}^{n \times n}$  is a fancy object called a (*non-commutative*) *algebra*. Informally, this is a vector space with an additional operation that lets us multiply two vectors together (which, if you look closely, isn't a feature of vector spaces normally). This is outside the scope of the course, but it may be interesting to you to know that such things exist and that these properties are not unique to  $\mathbb{R}^{n \times n}$ .

The proof of this theorem will require the properties of the dot product (recall Proposition 3.1.6).

*Proof.* For simplicity, we'll introduce some notation. For a matrix  $M$

- (this one is standard notation)  $M_{ij}$  denotes the  $(i, j)^{\text{th}}$  entry of  $M$ ,
- $\text{row}_i(M)$  denotes the  $i^{\text{th}}$  row of  $M$ , and
- $\text{col}_j(M)$  denotes the  $j^{\text{th}}$  column of  $M$ .

- (a) This is clunky to do fully. Instead we'll note that for  $A \in \mathbb{R}^{m \times n}$ ,  $B \in \mathbb{R}^{n \times p}$  and  $C \in \mathbb{R}^{p \times r}$ , both products  $(AB)C$  and  $A(BC)$  have the same size  $m \times r$ , so at least there's hope these will be equal as matrices. Instead of proving the rest of this part, we'll just do an example to demonstrate that it's true. [Add proof later maybe?]
- (b) Let  $A \in \mathbb{R}^{m \times n}$  and  $B, C \in \mathbb{R}^{n \times p}$ . Notice that  $A(B + C)$  and  $AB + AC$  have the same size, hence they are equal if they have the same corresponding elements.

$$\begin{aligned} (A(B + C))_{ij} &= \text{row}_i(A) \cdot \text{col}_j(B + C) \\ &= \text{row}_i(A) \cdot (\text{col}_j(B) + \text{col}_j(C)) \\ &= \text{row}_i(A) \cdot \text{col}_j(B) + \text{row}_i(A) \cdot \text{col}_j(C) = (AC)_{ij} + (BC)_{ij}. \end{aligned}$$

- (c) Let  $A, B \in \mathbb{R}^{m \times n}$  and  $C \in \mathbb{R}^{n \times p}$ . Notice that  $(A + B)C$  and  $AC + BC$  have the same size, hence they are equal if they have the same corresponding elements.

$$\begin{aligned} ((A + B)C)_{ij} &= \text{row}_i(A + B) \cdot \text{col}_j(C) \\ &= (\text{row}_i(A) + \text{row}_i(B)) \cdot \text{col}_j(C) \\ &= \text{row}_i(A) \cdot \text{col}_j(C) + \text{row}_i(B) \cdot \text{col}_j(C) = (AC)_{ij} + (BC)_{ij}. \end{aligned}$$

- (d) Let  $A \in \mathbb{R}^{m \times n}$ ,  $B \in \mathbb{R}^{n \times p}$ , and  $k \in \mathbb{R}$ . Notice that  $k(AB)$ ,  $(kA)B$  and  $A(kB)$  all have the same size  $m \times p$ , hence they are equal if they have the same corresponding elements.

$$\begin{aligned} (k(AB))_{ij} &= k(\text{row}_i(A) \cdot \text{col}_j(B)) \\ &= \text{row}_i(kA) \cdot \text{col}_j(B) = ((kA)B)_{ij} \\ &= \text{row}_i(A) \cdot \text{col}_j(kB) = (A(kB))_{ij} \end{aligned}$$

- (e) Let  $A \in \mathbb{R}^{m \times n}$ . Writing the  $m \times m$  identity matrix  $I_m = [\delta_{ij}]$  using the Kroenecker delta (c.f. page 13), we note that  $I_m A$  and  $A$  have the same size, hence they are equal if they have the same corresponding elements.

$$\begin{aligned} (I_m A)_{ij} &= \text{row}_i(I_m) \cdot \text{col}_j(A) \\ &= \delta_{i1}A_{1j} + \delta_{i2}A_{2j} + \cdots + \delta_{im}A_{mj} \\ &= \delta_{ii}A_{ij} && \text{(the only nonzero term)} \\ &= A_{ij} \end{aligned}$$

Similarly, for the  $n \times n$  identity matrix  $I_n$ ,

$$\begin{aligned} (AI_n)_{ij} &= \text{row}_i(A) \cdot \text{col}_j(I_n) \\ &= A_{i1}\delta_{1j} + A_{i2}\delta_{2j} + \cdots + A_{in}\delta_{nj} \\ &= A_{ij}\delta_{jj} && \text{(the only nonzero term)} \\ &= A_{ij} \end{aligned}$$

□

**Example 3.2.3.** Let  $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix}$ ,  $B = \begin{bmatrix} 3 & 2 \\ 1 & 0 \\ -1 & -2 \end{bmatrix}$ , and  $C = \begin{bmatrix} 1 & -2 \\ 3 & -4 \end{bmatrix}$ . Then

$$\begin{aligned} A(BC) &= \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \left( \begin{bmatrix} 3 & 2 \\ 1 & 0 \\ -1 & -2 \end{bmatrix} \begin{bmatrix} 1 & -2 \\ 3 & -4 \end{bmatrix} \right) \\ &= \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \left( \begin{bmatrix} 9 & -14 \\ 1 & -2 \\ -7 & 10 \end{bmatrix} \right) \\ &= \begin{bmatrix} -10 & 12 \\ -1 & -6 \end{bmatrix} \end{aligned}$$

and

$$\begin{aligned} (AB)C &= \left( \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 1 & 0 \\ -1 & -2 \end{bmatrix} \right) \begin{bmatrix} 1 & -2 \\ 3 & -4 \end{bmatrix} \\ &= \left( \begin{bmatrix} 2 & -4 \\ 11 & -4 \end{bmatrix} \right) \begin{bmatrix} 1 & -2 \\ 3 & -4 \end{bmatrix} \\ &= \begin{bmatrix} -10 & 12 \\ -1 & -6 \end{bmatrix} \end{aligned}$$

**Theorem 3.2.4** (*Poole Theorem 3.4 - Properties of the Transpose*). Let  $A$  and  $B$  be matrices (whose sizes are such that the indicated operations can be performed) and let  $k$  be a scalar. Then

- (a)  $(A^T)^T = A$
- (b)  $(kA)^T = k(A^T)$
- (c)  $(A^r)^T = (A^T)^r$  for all nonnegative integers  $r$ .
- (d)  $(A + B)^T = A^T + B^T$
- (e)  $(AB)^T = B^T A^T$

*Proof.* We use the same notation as in the proof of Theorem 3.2.2.

- (a) [Add proof later?]
- (b) [Add proof later?]
- (c) [Add proof later?]
- (d) [Add proof later?]
- (e) Let  $A \in \mathbb{R}^{m \times n}$  and  $B \in \mathbb{R}^{n \times p}$ . Note that  $(AB)^T$  and  $B^T A^T$  both have the same size, hence they are equal if their corresponding entries are equal.

$$\begin{aligned} ((AB)^T)_{ij} &= (AB)_{ji} \\ &= \text{row}_j(A) \cdot \text{col}_i(B) \\ &= \text{col}_j(A^T) \cdot \text{row}_i(B^T) \\ &= \text{row}_i(B^T) \cdot \text{col}_j(A^T) = B^T A^T. \end{aligned}$$

□

## Section 3.2 Exercises

- (a) A matrix  $A \in \mathbb{R}^{n \times n}$  is **idempotent** if  $A^2 = A$  and these are quite useful in statistics. Give an example of a  $2 \times 2$  matrix (not equal to  $I_2$  or  $O_{2 \times 2}$ ) that is idempotent. (HINT: This can be done with just 1's and 0's.)

(b) With the matrix that you found in part (a), now compute  $(I_2 - A)$  and  $(I_2 - A)^2$ . What can you say about  $(I_2 - A)$ ?
- A matrix  $A \in \mathbb{R}^{n \times n}$  is **block diagonal** if it is of the form

$$A = \begin{bmatrix} B_1 & & & \\ & B_2 & & \\ & & \ddots & \\ & & & B_k \end{bmatrix}$$

where each of these  $B_i$  “blocks” are square matrices of varying sizes (and all empty entries are assumed to be zero). Block diagonal matrices are very nice for computation. Compute the following product of block-diagonal matrices. Do you notice anything interesting?

$$\begin{bmatrix} 1 & 2 & 0 & 0 \\ 3 & 4 & 0 & 0 \\ 0 & 0 & 5 & 6 \\ 0 & 0 & 7 & 8 \end{bmatrix} \begin{bmatrix} 16 & 15 & 0 & 0 \\ 14 & 13 & 0 & 0 \\ 0 & 0 & 12 & 11 \\ 0 & 0 & 10 & 9 \end{bmatrix} = ?$$

## 2.1 Introduction to Linear Systems

**Definition.** A *linear equation* in the variables  $x_1, \dots, x_n$  is an equation that can be written in the form

$$a_1x_1 + \dots + a_nx_n = b$$

where  $a_1, \dots, a_n$  are real numbers called *coefficients* and  $b$  is a real number called the *constant term*. A *solution* of the equation is a vector  $[v_1, \dots, v_n]$  so that

$$a_1v_1 + \dots + a_nv_n = b.$$

**Example 2.1.1.**  $4x - y = 2$  is an example of a linear equation. And notice we can rearrange it as  $y = 4x - 2$ , which is the equation of a line (hence why we call these “linear”). The vector  $[1, 2]$  is a solution because

$$4(1) - (2) = 2.$$

In fact, for any real number  $t$ , the vector  $[t, 4t - 2]$  is a solution because

$$4(t) - (4t - 2) = 2.$$

This means there are infinitely many possible solutions.

**Definition.** The collection of all solutions to a linear equation is called the *solution set* of that equation.

**Example 2.1.2.**  $\sin\left(\frac{\pi}{82364423}\right)x + \sqrt{540.6464}y + z = e^{71}$  is a linear equation, because complicated as they are,  $\sin\left(\frac{\pi}{82364423}\right)$ ,  $\sqrt{540.6464}$ ,  $e^{71}$  are just real numbers.

**Example 2.1.3.**  $x + xy + y = 7$  is *not* a linear equation because of that  $xy$  term.

**Example 2.1.4.**  $x^2 + 3^y = 8$  is *not* a linear equation because of both the  $x^2$  and  $3^y$  terms.

**Definition.** A *system of linear equations* is a finite set of linear equations, each with the same variables (and probably different coefficients). A *solution* of a system of linear equations is a vector that is simultaneously a solution for each linear equation in the system. A *solution set* is the collection of all possible solutions to the system.

**Example 2.1.5.** The system

$$\begin{cases} 2x - y = 3 \\ x + 3y = 5 \end{cases}$$

has the vector  $[2, 1]$  as a solution; in fact, this is the only solution.

**So, why vectors?**

Suppose we make a matrix of the coefficients of the system (the *coefficient matrix*:

$$A = \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix},$$

We also consider the column vector of the indeterminates  $\mathbf{x} = [x, y]^T$  and the system's constants as a column vector  $\mathbf{b} = [3, 5]^T$ . Notice that the product  $A\mathbf{x}$  is another  $2 \times 1$  matrix (i.e. a column vector)

$$A\mathbf{x} = \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2x - y \\ x + 3y \end{bmatrix}.$$

What we notice is that the vectors  $A\mathbf{x}$  and  $\mathbf{b}$  are equal if and only if the system above is satisfied. Notice that when  $\mathbf{x} = [2, 1]^T$ ,

$$A\mathbf{x} = \begin{bmatrix} 2 & -1 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 5 \end{bmatrix}$$

so a solution to a system of linear equations corresponds to a (vector) solution to the matrix/vector equation

$$A\mathbf{x} = \mathbf{b}.$$

This also suggests that our weird definition of matrix multiplication doesn't totally appear out of thin air. It's not the full story, but that will come in a later section. One may also notice

**Definition.** A system of linear equations is called *consistent* if it has at least one solution, and *inconsistent* if it has no solutions.

*Fact.* A system of linear equations with real coefficients has either

- (a) a unique solution (consistent)
- (b) infinitely many solutions (consistent)
- (c) no solutions (inconsistent)

You can convince yourself of the above trichotomy by considering how it works for systems of equations with two variables (whose solution sets are graphically lines in the Cartesian plane). Two lines can either intersect in a single point (if they are transverse), intersect in infinitely many points (if they coincide), or no points (if they are parallel).

**Example 2.1.6.** The system in Example 2.1.5 is consistent and the solution is unique.

**Example 2.1.7.** The system

$$\begin{cases} x - y = 0 \\ 2x - 2y = 0 \end{cases}$$

is consistent. It has the solution  $[x, y] = [1, 1]$ , but this is not the only solution. For any real number  $t$ , the vector  $[t, t]$  is a solution, so there are infinitely many.

**Example 2.1.8.** The system

$$\begin{cases} x + y = 0 \\ x + y = 2 \end{cases}$$

has no solutions.

**Definition.** Two systems of linear equations are called *equivalent* if they have the same solution set.





## Section 2.1 Exercises

1. Find the augmented matrix for each of the following systems:

$$(a) \begin{cases} x_1 = 3 - x_2 + x_3 \\ x_2 = x_4 - x_3 \\ x_3 = x_1 + x_4 + 1 \end{cases}$$

$$(b) \begin{cases} u = v \\ v = w \end{cases}$$

2. Find a system of two linear equations in the variables  $x$  and  $y$  whose solution set is given by the parametric equations  $x = t$  and  $y = 5 - 7t$ .
3. If the following linear system is **inconsistent**, what can be said about the value of  $h$ ?

$$x \begin{bmatrix} 1 \\ 2 \end{bmatrix} + y \begin{bmatrix} 4 \\ h \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \end{bmatrix}$$

## 2.2 Direct Methods for Solving Linear Systems

That “triangular structure” of systems of linear equations we saw previously makes life so easy... We should probably give it a name.

**Definition.** A matrix is in *row echelon form* if it satisfies the following properties:

- Any rows consisting entirely of zeros are at the bottom
- In each nonzero row, the first nonzero entry (the *leading entry*) is in a column to the left of any leading entries below it.

*Remark.* For a given matrix, row echelon form is not at all unique.

**Example 2.2.1.** The following matrices are in row echelon form.

$$\begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix} \quad \begin{bmatrix} 0 & 1 & 2 & 3 & 4 & 5 \\ 0 & 0 & 6 & 7 & 8 & 9 \\ 0 & 0 & 0 & 0 & 10 & 11 \\ 0 & 0 & 0 & 0 & 0 & 5 \end{bmatrix}$$

**Example 2.2.2.** In Example 2.1.5 we saw that the system

$$\begin{cases} 2x - y = 3 \\ x + 3y = 5 \end{cases}$$

was consistent and had the unique solution  $[x, y] = [2, 1]$ . Consider now the following three systems of equations. The first system is obtained by merely swapping the equations. The second system is obtained by scaling the second equation. The third system is obtained by replacing the second equation with the sum of the first and second equations.

$$\begin{cases} x + 3y = 5 \\ 2x - y = 3 \end{cases} \quad \begin{cases} 2x - y = 3 \\ 100x + 300y = 500 \end{cases} \quad \begin{cases} 2x - y = 3 \\ 3x + 2y = 8 \end{cases}$$

All of these have the (unique) solution  $[x, y] = [2, 1]$ , and so they are all equivalent.

It turns out that this fact isn't specific to this system, but is generally true of any linear system: these three operations do not change the solution set of the system! The “elimination method” (which you may be familiar with from a previous algebra/precalculus class) uses this fact to solve systems of linear equations. If we think about what this is doing to the corresponding augmented matrices, we get what we call the *elementary row operations*

**Definition.** For a given matrix, *elementary row operations* are any of the following operations:

- Swapping Row  $i$  and Row  $j$  (denoted  $R_i \leftrightarrow R_j$ ).
- Multiplying Row  $i$  by a nonzero constant (denoted  $kR_i \mapsto R_i$ ).
- Adding (a multiple of) Row  $j$  to Row  $i$  (denoted  $R_i + kR_j \mapsto R_i$ ).

*Remark.* These operations are not specific to augmented matrices, but are true of any matrices. In fact, unless explicitly stated otherwise, you should probably not ever assume that a matrix is augmented.

**Example 2.2.3.** In the interest of being super explicit, let's consider the following system and look at the interplay between the "elimination method" operations and the elementary row operations. For simplicity,  $E_i$  will refer to the  $i^{\text{th}}$  equation in the linear system, and  $R_j$  will refer to the  $j^{\text{th}}$  row of the augmented matrix.

Initial system of linear equations.

$$\begin{cases} x - y - z = 2 \\ 3x - 3y + 2z = 16 \\ 2x - y + z = 9 \end{cases}$$

Subtract  $3E_1$  from  $E_2$ .

$$\begin{cases} x - y - z = 2 \\ 5z = 10 \\ 2x - y + z = 9 \end{cases}$$

Subtract  $2E_1$  from  $E_3$ .

$$\begin{cases} x - y - z = 2 \\ 5z = 10 \\ y + 3z = 5 \end{cases}$$

Swap  $E_2$  and  $E_3$ .

$$\begin{cases} x - y - z = 2 \\ y + 3z = 5 \\ 5z = 10 \end{cases}$$

Corresponding augmented matrix.

$$\left[ \begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 3 & -3 & 2 & 16 \\ 2 & -1 & 1 & 9 \end{array} \right]$$

Subtract  $3R_1$  from  $R_2$ .

$$R_2 - 3R_1 \rightarrow R_2 \left[ \begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 0 & 0 & 5 & 10 \\ 2 & -1 & 1 & 9 \end{array} \right]$$

Subtract  $2R_1$  from  $R_3$ .

$$R_3 - 2R_1 \rightarrow R_3 \left[ \begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 0 & 0 & 5 & 10 \\ 0 & 1 & 3 & 5 \end{array} \right]$$

Swap  $R_2$  and  $R_3$ .

$$R_2 \leftrightarrow R_3 \left[ \begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 5 & 10 \end{array} \right]$$

From here it is straightforward to back-substitute and solve the system, whose solution is  $[x, y, z] = [3, -1, 2]$ .

**Definition.** The process of putting a matrix into row echelon form via these elementary row operations is called *row reduction* and the entry chosen to be a leading entry is sometimes called a *pivot*.

*Remark.* Often times authors will use *row reduction* to mean putting a matrix into *reduced* row echelon form (which we will describe shortly).

The definition of row echelon form suggests an algorithm for row reducing.

1. Working left to right, find the first nonzero column in the matrix.
2. Among all of the rows with nonzero entries in this column, choose one and move it to Row 1.
3. Use elementary row operations to clear all other nonzero entries in this column (below Row 1).
4. Ignoring Row 1, find the next nonzero column in this matrix.
5. Among all of the rows below Row 1 with nonzero entries in this column, choose one and move it to Row 2.
6. Use elementary row operations to clear all other nonzero entries in this column (below Row 2).
7. Repeat this process until the matrix is in Row echelon form.

**Definition.** Two matrices  $A$  and  $B$  are *row equivalent* if there is a sequence of elementary row operations taking  $A$  to  $B$ .

If  $A$  and  $B$  are augmented matrices, then this definition just tells us that  $A$  and  $B$  represent equivalent linear systems.

**Theorem 2.2.4** (Poole Theorem 2.1). *Matrices  $A$  and  $B$  are row equivalent if and only if they can be row reduced to the same echelon form.*

**Definition.** *Gaussian elimination* is the process from the previous example:

1. Write a linear system as an augmented matrix
2. Row reduce
3. Use back-substitution to solve the system.

**Example 2.2.5.** Use Gaussian elimination to find the solution set for the given system

$$\begin{cases} x_1 - x_2 + x_3 + 4x_4 = 0 \\ 2x_1 + x_2 - x_3 + 2x_4 = 9 \end{cases}$$

We set up the augmented matrix and row-reduce

$$\left[ \begin{array}{cccc|c} 1 & -1 & 1 & 4 & 0 \\ 2 & 1 & -1 & 2 & 9 \end{array} \right] \xrightarrow{R_2 - 2R_1 \mapsto R_2} \left[ \begin{array}{cccc|c} 1 & -1 & 1 & 4 & 0 \\ 0 & 3 & -3 & -6 & 9 \end{array} \right]$$

The corresponding system has infinitely-many solutions.

$$\begin{cases} x_1 - x_2 + x_3 + 4x_4 = 0 \\ 3x_2 - 3x_3 - 6x_4 = 9 \end{cases}$$

We can solve for the *leading variables* ( $x_1$  and  $x_2$ ) and do back-substitution to simplify

$$\begin{cases} x_1 = x_2 - x_3 - 4x_4 \\ x_2 = 3 + x_3 + 2x_4 \end{cases} \longrightarrow \begin{cases} x_1 = 3 - 2x_4 \\ x_2 = 3 + x_3 + 2x_4 \end{cases}$$

This form tells us that  $x_3$  and  $x_4$  are our *free variables*. Replacing them with parameters  $s$  and  $t$  (respectively), the vector solution is

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 3 - 2t \\ 3 + s + 2t \\ s \\ t \end{bmatrix}$$

which we can write as a linear combination of vectors with coefficients 1,  $s$ , and  $t$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} 3 - 2t \\ 3 + s + 2t \\ s \\ t \end{bmatrix} = \begin{bmatrix} 3 \\ 3 \\ 0 \\ 0 \end{bmatrix} + s \begin{bmatrix} 0 \\ 1 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -2 \\ 2 \\ 0 \\ 1 \end{bmatrix}.$$

**Definition.** The *reduced row echelon form* of a matrix is essentially the same as the row echelon form with the following additional requirements

1. Each leading entry is 1.
2. Any entries above a *leading 1* are also 0.

**Example 2.2.6.** The following matrices are in reduced row echelon form.

$$\begin{bmatrix} 1 & 0 & 3 \\ 0 & 1 & 5 \\ 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \begin{bmatrix} 0 & 1 & 0 & 3 & 0 & 0 \\ 0 & 0 & 1 & 7 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

*Remark.* The reduced row echelon form of a matrix is unique.

**Example 2.2.7.** Continuing with the system from Example 2.2.3, we'll put the row echelon form matrix into reduced row echelon form.

Scale  $E_3$  by  $\frac{1}{5}$ .

$$\begin{cases} x - y - z = 2 \\ y + 3z = 5 \\ z = 2 \end{cases}$$

Add  $E_2$  to  $E_1$ .

$$\begin{cases} x + 2z = 7 \\ y + 3z = 5 \\ z = 2 \end{cases}$$

Subtract  $2E_3$  from  $E_1$ .

Also subtract  $3E_3$  from  $E_2$ .

$$\begin{cases} x = 3 \\ y = -1 \\ z = 2 \end{cases}$$

Scale  $R_3$  by  $\frac{1}{5}$ .

$$\begin{array}{ccc|c} 1 & -1 & -1 & 2 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 1 & 2 \end{array} \quad \frac{1}{5}R_3 \mapsto R_3$$

Add  $R_2$  to  $R_1$ .

$$\begin{array}{ccc|c} 1 & 0 & 2 & 7 \\ 0 & 1 & 3 & 5 \\ 0 & 0 & 1 & 2 \end{array} \quad R_1 + R_2 \mapsto R_1$$

Subtract  $2R_3$  from  $R_1$ .

Also subtract  $3R_3$  from  $R_2$ .

$$\begin{array}{ccc|c} 1 & 0 & 0 & 3 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 2 \end{array} \quad \begin{array}{l} R_1 - 2R_3 \mapsto R_1 \\ R_2 - 3R_3 \mapsto R_2 \end{array}$$

Notice then that the solution  $[3, -1, 2]^T$  is even easier to read off of the matrix.

**Definition.** *Gauss-Jordan Elimination* is the same as Gaussian-elimination, except using the reduced row echelon form:

1. Write a linear system as an augmented matrix.
2. Row reduce (to reduced row echelon form).
3. Solve the system in terms of any remaining free variables.

What we have seen is that both row echelon form and reduced row echelon form are useful in the same way, but both have pros and cons. Row echelon form isn't unique and, in the case of augmented matrices, it takes a little bit more work to solve the system at the end. Reduced row echelon form is unique and makes the solution at the end easier, but requires more steps initially.

**What else can we do with (reduced) row echelon form?**

**Example 2.2.8.** In Example 2.1.8, we stated that the system

$$\begin{cases} x + y = 0 \\ x + y = 2 \end{cases}$$

was inconsistent. Look at what happens when we set up the augmented matrix and row-reduce:

$$\left[ \begin{array}{cc|c} 1 & 1 & 0 \\ 1 & 1 & 2 \end{array} \right] \xrightarrow{R_2 - R_1 \mapsto R_2} \left[ \begin{array}{cc|c} 1 & 1 & 0 \\ 0 & 0 & 2 \end{array} \right]$$

That last row corresponds to the linear equation  $0 = 2$ , which is patently false. This means there can't possibly be a solution to the system, i.e., it is inconsistent. We state this observation as a proposition.

**Proposition 2.2.9.** *If  $A = [a_{ij}|b_i]$  is an augmented matrix representing a system of linear equations and its (reduced) row echelon form contains a row where  $a_{i1} = a_{i2} = \cdots = a_{im} = 0$  and  $b_m \neq 0$ , then the system is inconsistent.*

One might ask if we can say anything about a consistent system from its (reduced) row echelon form. To answer this, we first introduce the following definition.

**Definition.** The *rank* of a matrix  $A$  is the number of nonzero rows in its (reduced) row echelon form, and is denoted  $\text{Rank}(A)$ .

**Example 2.2.10.** The rank of the coefficient matrix in Example 2.1.8 is 1, and the rank of the coefficient matrix in Example 2.2.3 is 2.

**Theorem 2.2.11** (Poole Theorem 2.2 - The Rank Theorem). *If  $A$  is the coefficient matrix of a consistent system of linear equations with  $n$  variables, then*

$$n = \text{Rank}(A) + \text{number of free variables.}$$

*Remark.* It turns out this theorem is actually just a special interpretation of a much more powerful theorem called the “Rank-Nullity Theorem,” but that discussion will have to wait for a later section.

**Definition.** A system of linear equations,  $A\mathbf{x} = \mathbf{b}$  is *homogeneous* if  $\mathbf{b} = \mathbf{0}$ .

*Remark.* Homogeneous systems are nice because they ALWAYS have at least one solution (namely the *trivial solution*  $\mathbf{x} = \mathbf{0}$ ).

**Theorem 2.2.12.** *If  $A\mathbf{x} = \mathbf{0}$  is a homogeneous system of  $m$  linear equations and  $n$  variables, where  $m < n$ , then the system has infinitely many solutions.*

*Proof.* Since the system is homogeneous, it has at least one solution. Since  $\text{Rank}(A) \leq m$ , then by the Rank Theorem

$$\text{number of free variables} = n - \text{Rank}(A) \geq n - m > 0$$

and a nonzero number of free variables implies that there are infinitely-many solutions. □

**Example 2.2.13.** Use Gaussian-Jordan elimination to find the solution set for the given system

$$\begin{cases} x_1 - x_2 + 3x_3 + 4x_4 = 0 \\ x_1 + x_2 - x_3 - 2x_4 = 0 \end{cases}$$

Creating the augmented matrix and doing the corresponding row operations, we have

$$\begin{aligned} \left[ \begin{array}{cccc|c} 1 & -1 & 3 & 4 & 0 \\ 1 & 1 & -1 & -2 & 0 \end{array} \right] &\xrightarrow{R_2 - R_1 \mapsto R_2} \left[ \begin{array}{cccc|c} 1 & -1 & 3 & 4 & 0 \\ 0 & 2 & -4 & -6 & 0 \end{array} \right] \\ &\xrightarrow{\frac{1}{2}R_2 \mapsto R_2} \left[ \begin{array}{cccc|c} 1 & -1 & 3 & 4 & 0 \\ 0 & 1 & -2 & -3 & 0 \end{array} \right] \\ &\xrightarrow{R_1 + R_2 \mapsto R_1} \left[ \begin{array}{cccc|c} 1 & 0 & 1 & 1 & 0 \\ 0 & 1 & -2 & -3 & 0 \end{array} \right] \end{aligned}$$

From here, we can see that  $x_3$  and  $x_4$  are free variables, so letting  $x_3 = s$  and  $x_4 = t$ , we get that the solution is

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \end{bmatrix} = \begin{bmatrix} -s - t \\ 2s + 3t \\ s \\ t \end{bmatrix} = s \begin{bmatrix} -1 \\ 2 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -1 \\ 3 \\ 0 \\ 1 \end{bmatrix}$$

The way this last example differs from Example 2.2.5 is that we have exactly as many free variables as we have vectors in the linear combination (instead of also having the extra constant vector added on). This is more ideal because, with the usual vector operations, the collection of all of these solutions is actually a vector space! We will explore this idea a bit further in the next section.

## Section 2.2 Exercises

1. Find a square matrix  $A$  such that...

(a)  $A$  that swaps the rows of a  $2 \times n$  matrix, i.e.

$$A \begin{bmatrix} x_1 & x_2 & \cdots & x_n \\ y_1 & y_2 & \cdots & y_n \end{bmatrix} = \begin{bmatrix} y_1 & y_2 & \cdots & y_n \\ x_1 & x_2 & \cdots & x_n \end{bmatrix}.$$

(HINT:  $A$  should only have 1's and 0's for its entries.)

(b)  $A$  that swaps the first and second rows of a  $3 \times n$  matrix (but fixes the third row), i.e.

$$A \begin{bmatrix} x_1 & x_2 & \cdots & x_n \\ y_1 & y_2 & \cdots & y_n \\ z_1 & z_2 & \cdots & z_n \end{bmatrix} = \begin{bmatrix} y_1 & y_2 & \cdots & y_n \\ x_1 & x_2 & \cdots & x_n \\ z_1 & z_2 & \cdots & z_n \end{bmatrix}.$$

(HINT:  $A$  should only have 1's and 0's for its entries.)

(c)  $A$  swaps the first and third rows of a  $3 \times n$  matrix (but fixes the second row).

(d)  $A$  swaps the second and third rows of a  $3 \times n$  matrix (but fixes the first row).

2. As you may have guessed by now, all elementary row operations can be represented as multiplication (on the left) by a square matrix called an **elementary matrix**. See if you can figure out how the other two elementary row operations work. For a real numbers  $k$ , find a square matrix  $A$  such that...

(a)  $A$  scales the second row of a  $2 \times n$  matrix by  $k$  (but fixes the first row), i.e.

$$A \begin{bmatrix} x_1 & x_2 & \cdots & x_n \\ y_1 & y_2 & \cdots & y_n \end{bmatrix} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \\ ky_1 & ky_2 & \cdots & ky_n \end{bmatrix}.$$

(b)  $A$  scales the second row of a  $3 \times n$  matrix by  $k$  (but fixes the first and third rows), i.e.

$$A \begin{bmatrix} x_1 & x_2 & \cdots & x_n \\ y_1 & y_2 & \cdots & y_n \\ z_1 & z_2 & \cdots & z_n \end{bmatrix} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \\ ky_1 & ky_2 & \cdots & ky_n \\ z_1 & z_2 & \cdots & z_n \end{bmatrix}.$$

(c)  $A$  adds the second row to the first row of a  $2 \times n$  matrix (and leaves the second row fixed), i.e.

$$A \begin{bmatrix} x_1 & x_2 \\ y_1 & y_2 \end{bmatrix} = \begin{bmatrix} (x_1 + y_1) & (x_2 + y_2) \\ y_1 & y_2 \end{bmatrix}.$$

(d)  $A$  adds the first row to the second row of a  $3 \times n$  matrix (but fixes the first and third rows), i.e.

$$A \begin{bmatrix} x_1 & x_2 & \cdots & x_n \\ y_1 & y_2 & \cdots & y_n \\ z_1 & z_2 & \cdots & z_n \end{bmatrix} = \begin{bmatrix} x_1 & x_2 & \cdots & x_n \\ (x_1 + y_1) & (x_2 + y_2) & \cdots & (x_n + y_n) \\ z_1 & z_2 & \cdots & z_n \end{bmatrix}.$$

3. For each of the following, give an example of two matrices  $A$  and  $B$  which have the given properties.

(a)  $\text{Rank}(A) = \text{Rank}(B)$  but  $A$  and  $B$  are not row equivalent.



- (b)  $A$  and  $B$  have the same size, but are not row equivalent.
4. [The Rank Theorem] Suppose  $A$  is a  $4 \times 4$  matrix,  $\mathbf{b}$  is a vector in  $\mathbb{R}^4$ , and the system  $A\mathbf{x} = \mathbf{b}$  is consistent. In each of the following cases, describe the number of solutions to the system. If there are infinitely-many, state how many parameters are needed for the solution set.
- $\text{Rank}(A) = 4$
  - $\text{Rank}(A) = 3$
  - $\text{Rank}(A) = 1$
5. Let  $A = \begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix}$  and let  $X = \begin{bmatrix} x_1 & x_2 \\ x_3 & x_4 \end{bmatrix}$  be a matrix of indeterminates.
- Find an augmented matrix representing the system produced by the matrix equation  $AX = I_2$ .
  - Use Gaussian or Gauss-Jordan elimination to solve this system.
  - With the solution from part (b) (rewritten as the matrix  $X$ ), what do you get when you compute the matrix product  $XA$ ?
  - What is  $\text{Rank}(A)$ ?
6. Let  $B = \begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix}$  and let  $X = \begin{bmatrix} x_1 & x_2 \\ x_3 & x_3 \end{bmatrix}$  be a matrix of indeterminates.
- Find an augmented matrix representing the system produced by the matrix equation  $BX = I_2$ .
  - Is this system consistent?
  - What is  $\text{Rank}(B)$ ?
7. Let  $A = \begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix}$  be the same matrix from Problem 5.
- Find matrices  $E_1, E_2, E_3$  corresponding to each of the following row operations

$$E_1 A = E_1 \begin{bmatrix} 1 & 3 \\ 2 & 5 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 0 & -1 \end{bmatrix} \quad (R_2 - 2R_1 \mapsto R_2)$$

$$E_2 \begin{bmatrix} 1 & 3 \\ 0 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} \quad (-R_2 \mapsto R_2)$$

$$E_3 \begin{bmatrix} 1 & 3 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = I_2 \quad (R_1 - 3R_2 \mapsto R_1)$$

- Compute the matrix product  $E_3 E_2 E_1$ . Do you notice anything interesting?

## 2.3 Spanning Sets and Linear Independence

Suppose that  $A$  is an  $m \times n$  matrix,  $\mathbf{x}$  is a vector in  $\mathbb{R}^n$  and  $\mathbf{b}$  is a vector in  $\mathbb{R}^m$ . Writing the columns of  $A$  as vectors  $\mathbf{a}_i$ , recall that the system  $A\mathbf{x} = \mathbf{b}$  expands as a linear combination of the columns of  $A$ :

$$\begin{aligned} A\mathbf{x} &= \begin{bmatrix} \mathbf{a}_1 & \cdots & \mathbf{a}_n \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} \\ &= \begin{bmatrix} a_{11} & \cdots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \cdots & a_{mn} \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} a_{11}x_1 + \cdots + a_{1n}x_n \\ \vdots \\ a_{m1}x_1 + \cdots + a_{mn}x_n \end{bmatrix} = x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n = \mathbf{b} \end{aligned}$$

**Theorem 2.3.1** (Poole Theorem 2.4). *A system of linear equations  $A\mathbf{x} = \mathbf{b}$  is consistent if and only if  $\mathbf{b}$  is a linear combination of the columns of  $A$ .*

The number of solutions to the system also tells us how many ways we can make such a linear combination. If there is a unique solution, then there is exactly one way. If there are infinitely-many solutions, there are infinitely-many ways to make the linear combination, so it may be reasonable to ask more qualitative questions about the set of all possible linear combinations and study the space of linear combinations instead.

**Definition.** Given a set of vectors  $S = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$  in a vector space  $V$ , we define the *span* of  $\mathbf{v}_1, \dots, \mathbf{v}_k$  to be the set of all linear combinations of these vectors, and we write  $\text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_k)$  or  $\text{Span}(S)$ . If  $V = \text{Span}(S)$ , then we call  $S$  a *spanning set* for  $V$ .

With this definition, we can restate Theorem 2.3.1 as follows:

**Theorem 2.3.2.** *A system of linear equations  $A\mathbf{x} = \mathbf{b}$  is consistent if and only if  $\mathbf{b}$  is in  $\text{Span}(\mathbf{a}_1, \dots, \mathbf{a}_n)$  (where  $\mathbf{a}_i$  is the  $i^{\text{th}}$  column of  $A$ ).*


**Exercise 2.3.1.** If  $V$  is a (real) vector space and  $\mathbf{v}_1, \dots, \mathbf{v}_k$  is some collection of vectors in  $V$ , then the set  $\text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_k)$  is also (real) vector space.

**Example 2.3.3.** Let  $\mathbf{e}_1 = [1, 0]$ ,  $\mathbf{e}_2 = [0, 1]$  be the standard basis vectors for  $\mathbb{R}^2$ . By definition, an arbitrary vector in  $\text{Span}(\mathbf{e}_1, \mathbf{e}_2)$  is of the form

$$x\mathbf{e}_1 + y\mathbf{e}_2 = x[1, 0] + y[0, 1] = [x, y]$$

and so  $\text{Span}(\mathbf{e}_1, \mathbf{e}_2) = \mathbb{R}^2$ .

**Example 2.3.4.** Let  $\mathbf{e}_1 = [1, 0, 0]^T$  and  $\mathbf{e}_2 = [0, 1, 0]^T$  be standard basis vectors in  $\mathbb{R}^3$ .  $\text{Span}(\mathbf{e}_1, \mathbf{e}_2)$  is the collection of all vectors in  $\mathbb{R}^3$  of the form  $[x, y, 0]$ , which is just the  $xy$ -plane. This set does not span  $\mathbb{R}^3$ , however, because it is missing all vectors with a nonzero 3<sup>rd</sup> coordinate (i.e. the  $z$ -direction).



**Example 2.3.5.** For any two vectors  $\mathbf{u}, \mathbf{v}$  in  $\mathbb{R}^3$  (with  $\mathbf{u}$  not a scalar multiple of  $\mathbf{v}$ ), then  $\text{Span}(\mathbf{u}, \mathbf{v})$  is a plane through the origin in  $\mathbb{R}^3$ .

Suppose we can write one vector  $\mathbf{w}$  as a linear combination of  $\mathbf{u}$  and  $\mathbf{v}$ , say  $\mathbf{w} = a\mathbf{u} + b\mathbf{v}$ . Then  $\mathbf{w}$  “depends” on  $\mathbf{u}$  and  $\mathbf{v}$ . Clearly we can rewrite this as

$$\mathbf{w} = a\mathbf{u} + b\mathbf{v} \quad \implies \quad a\mathbf{u} + b\mathbf{v} - \mathbf{w} = \mathbf{0},$$

and so we introduce the following definition.

**Definition.** A set of vectors  $\mathbf{v}_1, \dots, \mathbf{v}_k$  in a vector space is *linearly dependent* if there are scalars  $a_1, \dots, a_k$  (not all zero) such that

$$a_1\mathbf{v}_1 + \dots + a_k\mathbf{v}_k = \mathbf{0}.$$

A set of vectors that is not linearly dependent is called *linearly independent*.

*Remark.* It is always true that the equation above holds if  $a_1 = \dots = a_k = 0$ , so linearly dependence says that there is some other collection of  $a_i$ 's for which the equation is also true. In this way, linear independence can be thought of as saying that the only way the above equation is true is if  $a_1 = \dots = a_k = 0$ .

**Example 2.3.6.** The set  $\mathbf{v}_1 = [1, 1]$ ,  $\mathbf{v}_2 = [2, 1]$  and  $\mathbf{v}_3 = [1, 2]$  is linearly dependent because  $\mathbf{v}_2 + \mathbf{v}_3 - 3\mathbf{v}_1 = \mathbf{0}$ .

It should be clear from the way we defined linear dependence that the idea is to capture when one vector can be written as a linear combination of the others. In the above example we can easily write  $\mathbf{v}_3 = \frac{1}{3}\mathbf{v}_2 + \frac{1}{3}\mathbf{v}_2$ . In fact, this is an equivalent characterization of linear dependence.

**Theorem 2.3.7** (Poole Theorem 2.5). *The set of vectors  $S = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$  in a vector space is linearly dependent if and only if at least one of the vectors can be written as a linear combination of the others.*

The proof of this fact is essentially exactly what happens in the example, so we provide it fully below. The hard part is that we have to prove two separate things (because the statement of the theorem is a “biconditional statement”).

*Proof.* If  $S$  is linearly dependent, then we can find scalars  $a_1, \dots, a_k$ , not all zero, so that

$$a_1 \mathbf{v}_1 + \cdots + a_k \mathbf{v}_k = \mathbf{0}.$$

Since one of the coefficients  $a_i \neq 0$ , then we can rearrange this as

$$\mathbf{v}_i = -\frac{a_1}{a_i} \mathbf{v}_1 - \cdots - \frac{a_{i-1}}{a_i} \mathbf{v}_{i-1} - \frac{a_{i+1}}{a_i} \mathbf{v}_{i+1} - \cdots - \frac{a_k}{a_i} \mathbf{v}_k.$$

Conversely, suppose  $\mathbf{v}_1$  is nonzero and is a linear combination of the remaining vectors in  $S$ . Then there are constants  $a_2, \dots, a_k$ , not all zero, for which

$$\mathbf{v}_1 = a_2 \mathbf{v}_2 + \cdots + a_k \mathbf{v}_k$$

which rearranges to

$$-\mathbf{v}_1 + a_2 \mathbf{v}_2 + \cdots + a_k \mathbf{v}_k = \mathbf{0}.$$

hence  $S$  is linearly dependent. □

*Remark.* The above theorem is actually a bit subtle. It doesn't say that *every* vector can be written as a linear combination of the others, just that there's at least one that can be written this way.

### How can we use matrices to help us determine linear (in)dependence?

Given a collection of vectors  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ , determining linear (in)dependence comes down to finding whether (or not) there exist nonzero real numbers  $x_1, \dots, x_n$  such that

$$x_1 \mathbf{v}_1 + \cdots + x_n \mathbf{v}_n = \mathbf{0}$$

and this equivalent to checking whether or not the following system has any nontrivial solutions (i.e. solutions other than the zero vector).

$$A\mathbf{x} = \begin{bmatrix} \mathbf{v}_1 & \cdots & \mathbf{v}_n \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = x_1 \mathbf{v}_1 + \cdots + x_n \mathbf{v}_n = \mathbf{0}.$$

**Theorem 2.3.8** (Poole Theorem 2.6). *Let  $\mathbf{v}_1, \dots, \mathbf{v}_m$  be vectors in  $\mathbb{R}^n$  and let  $A$  be the  $n \times m$  matrix with these vectors as its columns,  $A = [\mathbf{v}_1, \dots, \mathbf{v}_m]$ . Then that collection of vectors is linearly dependent if and only if the homogeneous system  $[A|\mathbf{0}]$  has at least one nontrivial solution.*

The following theorem is logically equivalent to the above, but is stated to make the connection between this system and linear independence completely clear.

**Theorem 2.3.9.** *Let  $\mathbf{v}_1, \dots, \mathbf{v}_m$  be vectors in  $\mathbb{R}^n$  and let  $A$  be the  $n \times m$  matrix with these vectors as its columns,  $A = [\mathbf{v}_1, \dots, \mathbf{v}_m]$ . Then that collection of vectors is linearly independent if and only if the homogeneous system  $[A|\mathbf{0}]$  has no nontrivial solutions.*

**Example 2.3.10.** Consider the column vectors  $\mathbf{v}_1 = [0, 1, 2]^T$ ,  $\mathbf{v}_2 = [2, 1, 3]^T$ ,  $\mathbf{v}_3 = [2, 0, 2]^T$ . We can check for linear (in)dependence by row reducing  $[\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3 | \mathbf{0}]$  and checking for consistency.

$$\begin{aligned}
 [\mathbf{v}_1 \ \mathbf{v}_2 \ \mathbf{v}_3 | \mathbf{0}] &= \left[ \begin{array}{ccc|c} 0 & 2 & 2 & 0 \\ 1 & 1 & 0 & 0 \\ 2 & 3 & 2 & 0 \end{array} \right] \longrightarrow \begin{array}{c} \\ R_1 \leftrightarrow R_2 \\ \\ \end{array} \left[ \begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 2 & 2 & 0 \\ 2 & 3 & 2 & 0 \end{array} \right] \\
 &\longrightarrow \begin{array}{c} \\ \\ \frac{1}{2}R_2 \mapsto R_2 \\ R_3 - 2R_1 \mapsto R_3 \end{array} \left[ \begin{array}{ccc|c} 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 1 & 2 & 0 \end{array} \right] \\
 &\longrightarrow \begin{array}{c} R_1 - R_2 \mapsto R_1 \\ \\ R_3 - R_2 \mapsto R_3 \end{array} \left[ \begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right] \\
 &\longrightarrow \begin{array}{c} R_1 + R_3 \mapsto R_1 \\ R_2 - R_3 \mapsto R_2 \\ \\ \end{array} \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{array} \right]
 \end{aligned}$$

The system thus has no nontrivial solutions (because there are no free variables). By Theorem 2.3.9, the set  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  is linearly independent.

**Example 2.3.11.** Consider the column vectors  $\mathbf{v}_1 = [0, 1, 2]^T$ ,  $\mathbf{v}_2 = [2, 1, 3]^T$ ,  $\mathbf{v}_3 = [2, 0, 1]^T$ . We can check linear dependence by using Gaussian (or Gauss-Jordan) Elimination to see if  $[\mathbf{v}_1 \ \mathbf{v}_2 \ \mathbf{v}_3 | \mathbf{0}]$  is consistent.

$$\left[ \begin{array}{ccc|c} 0 & 2 & 2 & 0 \\ 1 & 1 & 0 & 0 \\ 2 & 3 & 1 & 0 \end{array} \right] \xrightarrow{\text{RREF}} \left[ \begin{array}{ccc|c} 1 & 0 & -1 & 0 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

It's an exercise to the reader to check that the reduced row echelon form is correct. The system thus has no nontrivial solutions. By Theorem 2.3.8 the set  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  is linearly dependent. Of course, this isn't hard to see, as  $\mathbf{v}_2 = \mathbf{v}_1 + \mathbf{v}_3$ .

**Example 2.3.12.** Let  $\mathbf{v}_1 = [1, 1]^T$ ,  $\mathbf{v}_2 = [1, -1]^T$ ,  $\mathbf{v}_3 = [x, y]^T$  be column vectors in  $\mathbb{R}^2$ . Using the same method from Theorem 2.3.8 we can check for linear dependence via Gaussian (or Gauss-Jordan) elimination:

$$[\mathbf{v}_1 \ \mathbf{v}_2 \ \mathbf{v}_3 | \mathbf{0}] = \left[ \begin{array}{ccc|c} 1 & 1 & x & 0 \\ 1 & -1 & y & 0 \end{array} \right] \xrightarrow{\text{RREF}} \left[ \begin{array}{ccc|c} 1 & 0 & \frac{x-y}{2} & 0 \\ 0 & 1 & \frac{x+y}{2} & 0 \end{array} \right]$$

It's an exercise to the reader to check that the reduced row echelon form is correct. The system has nontrivial solutions for every nonzero vector  $[x, y]^T$ , so by Theorem 2.3.8 the set  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  is linearly dependent.

Of course, three nonzero vectors in  $\mathbb{R}^2$  being linearly dependent doesn't sound too unreasonable - the corresponding linear system must have rank at most 2 and thus at least one free variable. We deduce from this the following result:

**Theorem 2.3.13** (*Poole Theorem 2.8*). *Any set of  $m$  vectors in  $\mathbb{R}^n$  is linearly dependent if  $m > n$ .*

### Why is linear independence important?

It all comes down to uniqueness. If  $\mathbf{v}_1, \dots, \mathbf{v}_n$  are linearly independent and  $\mathbf{b} \in \text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_n)$ , then linear independence of  $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$  tells us that the linear combination

$$\mathbf{b} = c_1\mathbf{v}_1 + \cdots + c_n\mathbf{v}_n$$

is actually the unique one representing  $\mathbf{b}$ . This is great because it means that if two people can agree on a particular (ordered) linearly independent set of vectors, then we can make a vector of the coefficients  $[c_1, \dots, c_n]^T$  that unambiguously represents the vector  $\mathbf{b}$ . Contrast this with the following example:

**Example 2.3.14.** Consider the vectors  $\mathbf{v}_1 = [1, 0]^T$ ,  $\mathbf{v}_2 = [0, 1]^T$ ,  $\mathbf{v}_3 = [1, 1]^T$ . Clearly, the set  $\{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$  is linearly dependent. So we can write  $\mathbf{b} = [3, 1]^T$  in multiple ways, say,

$$\mathbf{b} = 3\mathbf{v}_1 + 1\mathbf{v}_2 + 0\mathbf{v}_3 = 2\mathbf{v}_1 + 0\mathbf{v}_2 + 1\mathbf{v}_3 = 0\mathbf{v}_1 - 2\mathbf{v}_2 + 3\mathbf{v}_3$$

and if we tried to refer to  $\mathbf{b}$  as just a vector of its coefficients, we could have that  $[3, 1, 0]^T$ ,  $[0, 2, 1]^T$ , and  $[0, -2, 3]^T$  all represent the same vector.

This particular idea will become especially important when we discuss bases for a vector space, but there are also several other ideas that we will see are reliant upon the notion of linear independence.

## Section 2.3 Exercises

1. Give an example of three vectors  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^2$  for which  $\{\mathbf{u}, \mathbf{v}, \mathbf{w}\}$  is a linearly dependent set, but  $\mathbf{u}$  cannot be written as a linear combination of  $\mathbf{v}$  and  $\mathbf{w}$ .
2. (a) Let  $\mathbf{v}_1 = [1, 1]^T$  and  $\mathbf{v}_2 = [1, -1]^T$  be (column) vectors in  $\mathbb{R}^2$ . Show that  $\mathbb{R}^2 = \text{Span}(\mathbf{v}_1, \mathbf{v}_2)$  by showing that, for any arbitrary vector  $\mathbf{w} = [a, b]^T$ , the system  $[\mathbf{v}_1 \ \mathbf{v}_2 | \mathbf{w}]$  is consistent.  
(b) Let  $\mathbf{v}_1 = [1, 0, -1]^T$ ,  $\mathbf{v}_2 = [-1, 1, 0]^T$ , and  $\mathbf{v}_3 = [0, -1, 1]^T$  be (column) vectors in  $\mathbb{R}^3$ . Show that  $\mathbb{R}^3 \neq \text{Span}(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$  and geometrically describe  $\text{Span}(\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3)$ .
3. The following question appeared on the Fall 2016 common time final exam. Determine which answer is correct and provide an explanation for why each of the remaining three is incorrect.

Determine whether or not  $V$  spans  $\mathbb{R}^2$  and choose a correct justification:

$$V = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}, \quad \text{where } \mathbf{v}_1 = \begin{bmatrix} 3 \\ -1 \end{bmatrix}, \quad \mathbf{v}_2 = \begin{bmatrix} 2 \\ 7 \end{bmatrix}, \quad \mathbf{v}_3 = \begin{bmatrix} 6 \\ 21 \end{bmatrix}$$

- (a)  $V$  spans  $\mathbb{R}^2$  because its echelon form contains a leading entry in every column.
- (b)  $V$  spans  $\mathbb{R}^2$  because  $x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + x_3\mathbf{v}_3 = \mathbf{b}$  is consistent for all  $\mathbf{b} \in \mathbb{R}^2$ .
- (c)  $V$  does not span  $\mathbb{R}^2$  because  $\mathbf{v}_3$  is a scalar multiple of  $\mathbf{v}_2$ .
- (d)  $V$  does not span  $\mathbb{R}^2$  because a set of three vectors can only span  $\mathbb{R}^3$ .

### 3.3 The Inverse of a Matrix

Motivation: If  $a, b \in \mathbb{R}$  and  $x$  is some unknown and we wanted to solve for  $x$  in the equation

$$ax = b,$$

we would do so multiplying both sides by  $a^{-1} = \frac{1}{a}$  to get that  $x = a^{-1}b$ . We'd like to be able to do this same thing for the system of linear equations

$$A\mathbf{x} = \mathbf{b}$$

where  $A \in \mathbb{R}^{n \times n}$  and  $\mathbf{b} \in \mathbb{R}^n$ . But alas, we don't have a notion of division of matrices.

Let's think – if  $a \in \mathbb{R}$  is some nonzero number, then  $a^{-1}$  is just some other real number for which  $aa^{-1} = a^{-1}a = 1$ . Since the  $n \times n$  identity matrix  $I_n$  plays the role of 1, multiplicatively, then the natural way to define the matrix we desire is

**Definition.** For an nonzero  $n \times n$  matrix  $A$ , the *inverse of  $A$* , denoted  $A^{-1}$ , is the  $n \times n$  matrix satisfying

$$AA^{-1} = A^{-1}A = I_n.$$

If the inverse exists, we say that  $A$  is *invertible*.

*Fact.* Not every nonzero matrix is invertible, and we'll devote the latter half of this section to exploring when a matrix is invertible.

*Remark.* We only define inverses for square matrices. You will explore what happens in your homework for non-square matrices.

**Theorem 3.3.1** (Poole Theorem 3.6). *The inverse is unique.*

*Proof.* Suppose  $X, Y$  are both inverses of  $A$ . Then

$$X = X(I_n) = X(AY) = (XA)Y = (I_n)Y = Y.$$

□

**Theorem 3.3.2** (Poole Theorem 3.9). *If  $A, B \in \mathbb{R}^{n \times n}$  are invertible and  $c \in \mathbb{R}$  is some nonzero scalar, then*

- a.  $A^{-1}$  is invertible and  $(A^{-1})^{-1} = A$ .
- b.  $(cA)^{-1} = \frac{1}{c}A^{-1}$
- c.  $(AB)^{-1} = B^{-1}A^{-1}$
- d.  $A^T$  is invertible and  $(A^T)^{-1} = (A^{-1})^T$
- e.  $A^n$  is invertible for all positive integers  $n$  and  $(A^k)^{-1} = (A^{-1})^k$

*Proof.* Since inverses are unique, and matrices are invertible if their inverses exist, then each of these is essentially proven by merely checking that the multiplication checks out.

- a.  $(A^{-1})(A) = I_n$
- b.  $(cA)(\frac{1}{c}A^{-1}) = \frac{c}{c}AA^{-1} = I_n$ .



- c.  $(AB)(B^{-1}A^{-1}) = AI_nA^{-1} = AA^{-1} = I_n$ .
- d.  $A^T(A^{-1})^T = (A^{-1}A)^T = I_n^T = I_n$
- e.  $(A^k)(A^{-1})^k = \underbrace{A \cdots A}_k \underbrace{A^{-1} \cdots A^{-1}}_k = \underbrace{A \cdots A}_{k-1} I_n \underbrace{A^{-1} \cdots A^{-1}}_{k-1} = \cdots = I_n$

□

*Remark.* Because of the above theorem, some will use the notation  $A^{-n}$  (for  $n$  a positive integer) and  $A^{-T}$  (the transpose) to mean the obvious things:

$$A^{-n} = (A^{-1})^n = (A^n)^{-1}$$

$$A^{-T} = (A^T)^{-1} = (A^{-1})^T$$

**Suppose  $A$  is invertible. How do we find  $A^{-1}$ ?**

Well, you secretly already did this on your last homework assignment \*cue nefarious laughter\*. If  $A = [a_{ij}]$  is given and  $X = [x_{ij}]$  is a matrix of indeterminates, The matrix equation  $AX = I_n$  yields a linear system (with  $n^2$  equations and  $n^2$  unknowns)

$$\begin{cases} \sum_{k=1}^n a_{1k}x_{k1} & = 1 \\ \sum_{k=1}^n a_{1k}x_{k2} & = 0 \\ & \vdots \\ \sum_{k=1}^n a_{ik}x_{kj} & = \delta_{ij} \quad (\text{the Kroenecker delta}) \\ & \vdots \\ \sum_{k=1}^n a_{nk}x_{kn} & = 1 \end{cases}$$

and you can use standard techniques to solve this system.

**Example 3.3.3.** Find  $A^{-1}$  given  $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ .

Let  $X = \begin{bmatrix} x_1 & x_2 \\ x_3 & x_4 \end{bmatrix}$  be a matrix of indeterminates. Then the matrix equation

$$AX = I_2$$

$$\begin{bmatrix} x_1 + 2x_3 & x_2 + 2x_4 \\ 3x_1 + 4x_3 & 3x_2 + 4x_4 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

yields the system

$$\begin{cases} x_1 & + & 2x_3 & & = & 1 \\ & x_2 & & + & 2x_4 & = & 0 \\ 3x_1 & & + & 4x_3 & & = & 0 \\ & 3x_2 & & + & 4x_4 & = & 1 \end{cases}$$

And solving this in the usual way, we get that there is a unique solution  $x_1 = -2$ ,  $x_2 = 1$ ,  $x_3 = \frac{3}{2}$ ,  $x_4 = -\frac{1}{2}$ . So

$$A^{-1} = \begin{bmatrix} -2 & 1 \\ \frac{3}{2} & -\frac{1}{2} \end{bmatrix}.$$

While this technique does work, it seems to get unwieldy pretty quickly because the system size is huge. The other way to solve for inverses (which was also on your homework), was via elementary matrices.

**Definition.** An *elementary matrix* is a matrix obtained by performing an elementary row operation on the identity matrix.

The other way to think about it is that, given a matrix  $A \in \mathbb{R}^{m \times n}$ , an elementary matrix  $E \in \mathbb{R}^{m \times m}$  is one for which the product  $EA$  has the same effect as doing an elementary row operation on  $A$ .

One can then get the inverse as the product of all of the elementary matrices.

**Example 3.3.4.** Given  $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$ . Use elementary matrices to compute  $A^{-1}$ .

In the following string of equalities, we'll denote the row reduction on the left-hand side and the corresponding product by elementary matrices on the right-hand side.

$$\begin{aligned} \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix} &= A \\ R_2 - 3R_1 \mapsto R_2 \quad \begin{bmatrix} 1 & 2 \\ 0 & -2 \end{bmatrix} &= \begin{bmatrix} 1 & 0 \\ -3 & 1 \end{bmatrix} A \\ \frac{-1}{2} R_2 \mapsto R_2 \quad \begin{bmatrix} 1 & 2 \\ 0 & 1 \end{bmatrix} &= \begin{bmatrix} 1 & 0 \\ 0 & -\frac{1}{2} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -3 & 1 \end{bmatrix} A \\ R_1 - 2R_2 \mapsto R_1 \quad \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} &= \begin{bmatrix} 1 & -2 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -\frac{1}{2} \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -3 & 1 \end{bmatrix} A \\ I_2 &= \begin{bmatrix} -2 & 1 \\ \frac{3}{2} & -\frac{1}{2} \end{bmatrix} A \end{aligned}$$

so again we get that  $A^{-1} = \begin{bmatrix} -2 & 1 \\ \frac{3}{2} & -\frac{1}{2} \end{bmatrix}$ .

If we write the elementary matrices above as  $E_1, E_2, E_3$ , respectively, (so that  $A^{-1} = E_3E_2E_1$ ) then applying them to the matrix equation  $AX = I$  looks something like the following

$$\begin{aligned} AX &= I \\ E_1AX &= E_1 \\ E_2E_1AX &= E_2E_1 \\ E_3E_2E_1AX &= E_3E_2E_1 \\ IX &= E_3E_2E_1 \end{aligned}$$

What this tells us is that computing the inverse can just be done with row operations on an augmented matrix  $[A|I]$  until we get  $[I|A^{-1}]$  (your book calls this the *Gauss-Jordan Method*).

**Example 3.3.5.** With  $A$  as before

$$\begin{aligned} & \left[ \begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 3 & 4 & 0 & 1 \end{array} \right] = [A \mid I] \\ & \xrightarrow{R_2 - 3R_1 \mapsto R_2} \left[ \begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 0 & -2 & -3 & 1 \end{array} \right] \\ & \xrightarrow{-\frac{1}{2}R_2 \mapsto R_2} \left[ \begin{array}{cc|cc} 1 & 2 & 1 & 0 \\ 0 & 1 & \frac{3}{2} & -\frac{1}{2} \end{array} \right] \\ & \xrightarrow{R_1 - 2R_2 \mapsto R_1} \left[ \begin{array}{cc|cc} 1 & 0 & -2 & 1 \\ 0 & 1 & \frac{3}{2} & -\frac{1}{2} \end{array} \right] = [I \mid A^{-1}] \end{aligned}$$

One can, of course, do the above process for generic  $2 \times 2$  matrices, which yields the following result.

**Theorem 3.3.6** (Poole thm 3.8). *If  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ , then the inverse is given by  $A^{-1} = \frac{1}{ad-bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$ , provided  $ad - bc \neq 0$ .*

*Proof.* Using the Gauss-Jordan method above,

$$\begin{aligned} & \left[ \begin{array}{cc|cc} a & b & 1 & 0 \\ c & d & 0 & 1 \end{array} \right] \xrightarrow{\frac{1}{a}R_1 \mapsto R_1} \left[ \begin{array}{cc|cc} 1 & \frac{b}{a} & \frac{1}{a} & 0 \\ c & d & 0 & 1 \end{array} \right] \\ & \xrightarrow{R_2 - cR_1 \mapsto R_2} \left[ \begin{array}{cc|cc} 1 & \frac{b}{a} & \frac{1}{a} & 0 \\ 0 & \frac{ad-bc}{a} & -\frac{c}{a} & 1 \end{array} \right] \\ & \xrightarrow{\frac{a}{ad-bc}R_2 \mapsto R_2} \left[ \begin{array}{cc|cc} 1 & \frac{b}{a} & \frac{1}{a} & 0 \\ 0 & 1 & -\frac{c}{ad-bc} & \frac{a}{ad-bc} \end{array} \right] \\ & \xrightarrow{R_1 - \frac{b}{a}R_2 \mapsto R_1} \left[ \begin{array}{cc|cc} 1 & 0 & \frac{d}{ad-bc} & -\frac{b}{ad-bc} \\ 0 & 1 & -\frac{c}{ad-bc} & \frac{a}{ad-bc} \end{array} \right] \end{aligned}$$

□

You could do this same system-solving process for larger matrices, but the formulas are significantly worse.

**How does this help us solve our system?**

If  $A$  is invertible and  $A\mathbf{x} = \mathbf{b}$  then, by design, it should have the solution  $\mathbf{x} = A^{-1}\mathbf{b}$ . Moreover, since  $A^{-1}$  is unique, we expect this solution to be unique.

**Theorem 3.3.7** (Poole Theorem 3.7). *If  $A$  is an invertible  $n \times n$  matrix, then for every  $\mathbf{b} \in \mathbb{R}^n$ , the linear system  $A\mathbf{x} = \mathbf{b}$  is consistent and has the unique solution  $\mathbf{x} = A^{-1}\mathbf{b}$ .*

**Example 3.3.8.** Solve the system  $A\mathbf{x} = \mathbf{b}$  given  $A = \begin{bmatrix} 1 & 4 \\ 3 & 13 \end{bmatrix}$  and  $\mathbf{b} = \begin{bmatrix} 1 \\ 7 \end{bmatrix}$ .

We could solve this the old way, or we can try our nifty new method. We quickly deduce that  $A^{-1}$  is given by

$$A^{-1} = \begin{bmatrix} 13 & -4 \\ -3 & 1 \end{bmatrix}$$

(which can be seen either by appealing to Theorem 3.3.6 or using the Gauss-Jordan Method). Hence the solution is

$$\mathbf{x} = A^{-1}\mathbf{b} = \begin{bmatrix} 13 & -4 \\ -3 & 1 \end{bmatrix} \begin{bmatrix} 1 \\ 7 \end{bmatrix} = \begin{bmatrix} -15 \\ 4 \end{bmatrix}.$$

**Example 3.3.9.** Solve the system

$$\begin{cases} x + y = 1 \\ -x - y = -1 \end{cases}$$

Notice that this system is equivalent to the system  $\{x + y = 1\}$ , which has infinitely-many solutions. Notice also that the coefficient matrix for this system is

$$A = \begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$$

which isn't invertible (because otherwise, attempting to apply Theorem 3.3.6 we would be dividing by 0).

**Okay, so tell me, when is  $A$  invertible?**

Putting it all together, we can wrap it up into the following theorem

**Theorem 3.3.10** (Poole Theorem 3.12 - The Fundamental Theorem of Invertible Matrices: Pt I). *Let  $A \in \mathbb{R}^{n \times n}$ . The following are equivalent:*

- a.  $A$  is invertible.*
- b.  $A$  is row equivalent to  $I_n$  (i.e. its reduced row echelon form is  $I_n$ ).*
- c.  $A$  is the product of elementary matrices.*
- d.  $A\mathbf{x} = \mathbf{b}$  has a unique solution for every  $\mathbf{b} \in \mathbb{R}^n$ .*
- e.  $A\mathbf{x} = \mathbf{0}$  has only the trivial solution.*
- f. The columns of  $A$  are linearly independent.*

For part (c) above, we saw that it worked in one particular example. In fact, it's true for all  $\mathbf{b}$  because, if  $A$  is row equivalent to the identity, then it will never have a row of all 0's in its row-reduced form (which is precisely what happened in the cases that  $A\mathbf{x} = \mathbf{b}$  had no solutions or infinitely many solutions).

### Section 3.3 Exercises

1. **Why do we only define inverses for square matrices?** Suppose  $A$  is an  $m \times n$  matrix (with  $m \neq n$ ) and we wanted to try to find some matrix  $X$  so that  $XA = AX = I$  (the identity matrix).
  - (a) If  $XA = I$ , then what size must  $X$  have?
  - (b) If  $AX = I$ , then what size must  $X$  have?
  - (c) What can you conclude about the existence such an  $X$ ?
2. For a fixed matrix  $A$ , a **left inverse** is a matrix  $X$  so that  $XA = I$ , and a **right inverse** is a matrix  $Y$  so that  $AY = I$  (but we don't require  $XA = AY$ ).

- (a) Let  $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{bmatrix}$ , and let  $X$  and  $Y$  be  $3 \times 2$  matrices of indeterminates:

$$X = \begin{bmatrix} x_{11} & x_{12} \\ x_{21} & x_{22} \\ x_{31} & x_{32} \end{bmatrix} \quad \text{and} \quad Y = \begin{bmatrix} y_{11} & y_{12} \\ y_{21} & y_{22} \\ y_{31} & y_{32} \end{bmatrix}.$$

- i. How many solutions are there to  $XA = I_3$
  - ii. How many solutions are there to  $AY = I_2$ ?
  - iii. If  $A$  has size  $m \times n$  with  $m < n$ , what do you expect to be true about left and right inverses for  $A$ ?
- (b) Now let  $A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 1 \end{bmatrix}$ , and let  $X$  and  $Y$  be  $2 \times 3$  matrices of indeterminates:

$$X = \begin{bmatrix} x_{11} & x_{12} & x_{13} \\ x_{21} & x_{22} & x_{23} \end{bmatrix} \quad \text{and} \quad Y = \begin{bmatrix} y_{11} & y_{12} & y_{13} \\ y_{21} & y_{22} & y_{23} \end{bmatrix}.$$

- i. How many solutions are there to  $XA = I_2$
  - ii. How many solutions are there to  $AY = I_3$ ?
  - iii. If  $A$  is  $m \times n$  with  $m > n$ , what do you expect to be true about the existence of left and right inverses for  $A$ ?
3. Give an example of  $2 \times 2$  matrices  $A, B$  such that  $A, B$ , and  $(A + B)$  are all invertible, but  $(A + B)^{-1} \neq A^{-1} + B^{-1}$
  4. Compute the inverse of the block-diagonal matrix  $A$ . What do you notice about  $A^{-1}$ ?

$$A = \begin{bmatrix} 2 & 1 & 0 & 0 \\ 5 & 3 & 0 & 0 \\ 0 & 0 & 1 & 2 \\ 0 & 0 & 3 & 4 \end{bmatrix}$$

### 3.4 Subspaces, Basis, Dimension, and Rank

We've thought about solution sets as spans of vectors and also, alternatively, as lines and planes in 3-dimensional space. Now we'll formalize these ideas so that we can talk about these things in more generality. Recall the definition of a real vector space:

**Definition.** A (*real*) *vector space*  $V$  is a set of objects (called *vectors*) with two operations *vector addition* (denoted  $+$ ) and *scalar multiplication* (no symbol) satisfying the following properties: For all vectors  $\mathbf{u}, \mathbf{v}, \mathbf{w}$  and real numbers  $a, b$  (called *scalars*),

- (a)  $\mathbf{u} + \mathbf{v}$  is in  $V$  [closure of addition]
- (b)  $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$  [commutativity of addition]
- (c)  $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$  [associativity of  $+$ ]
- (d) There is some vector  $\mathbf{0}$ , called the *zero vector*, [additive identity]  
so that  $\mathbf{u} + \mathbf{0} = \mathbf{u}$  for all vectors  $\mathbf{u}$ .
- (e) For each  $\mathbf{u}$  in  $V$ , there is some vector  $-\mathbf{u}$  for [additive inverse]  
which  $\mathbf{u} + (-\mathbf{u}) = \mathbf{0}$ .
- (f)  $a\mathbf{u}$  is in  $V$  [closure of scalar mult.]
- (g)  $a(\mathbf{u} + \mathbf{v}) = a\mathbf{u} + a\mathbf{v}$  [distributivity]
- (h)  $(a + b)\mathbf{u} = a\mathbf{u} + b\mathbf{u}$  [distributivity]
- (i)  $(ab)\mathbf{u} = a(b\mathbf{u})$  [associativity of scalar mult.]
- (j)  $1\mathbf{u} = \mathbf{u}$  [scalar mult. identity]

With this in mind, we introduce the following definition:

**Definition.** Let  $V$  be a vector space and let  $W$  be a subset of vectors in  $V$ . We say that  $W$  is a *subspace* of  $V$  if it is also a vector space (with the same vector addition/scalar multiplication operations).

In order to check that a set of vectors is a subspace, one would have to check all of the axioms of the vector space definition – eww. Instead, here is an equivalent characterization of a subspace (note: this is typically a theorem in most textbooks, but your book presents it as the definition).

**Definition.** Let  $V$  be a vector space and let  $W$  be a subset of vectors in  $V$ .  $W$  is a *subspace* of  $V$  if it has the following properties:

1.  $\mathbf{0}$  is in  $W$  (where  $\mathbf{0}$  is the same zero vector in  $V$ ).
2. If  $\mathbf{u}, \mathbf{v} \in W$ , then  $\mathbf{u} + \mathbf{v} \in W$ . [closure of addition]
3. If  $\mathbf{u} \in W$  and  $k \in \mathbb{R}$  is a scalar, then  $k\mathbf{u} \in W$ . [closure of scalar multiplication]

(where vector addition and scalar multiplication in  $W$  are the same operations for  $V$ ).

**Example 3.4.1.** Every vector space  $V$  is a subspace of itself.

**Example 3.4.2.** For any vector space  $V$ , the set  $\{\mathbf{0}\}$  is a subspace of  $V$  (sometimes called the *trivial subspace*).

**Example 3.4.3.** Let  $W$  be the set of all vectors in  $\mathbb{R}^3$  of the form  $[x, y, 0]$ . Then  $W$  is a subspace of  $\mathbb{R}^3$  because

1.  $[0, 0, 0] \in W$
2.  $[x_1, y_1, 0] + [x_2, y_2, 0] = [x_1 + x_2, y_1 + y_2, 0] \in W$
3.  $k[x, y, 0] = [kx, ky, 0] \in W$

**Example 3.4.4.** Let  $W$  be the set of all vectors in  $\mathbb{R}^3$  of the form  $[x, y, 1]$ . Then  $W$  is not a subspace of  $\mathbb{R}^3$  because it fails property 1 in the definition: the zero vector  $[0, 0, 0]$  does not have the form  $[x, y, 1]$ . (It actually fails the other two properties as well, but this is left as an exercise to the reader).

**Theorem 3.4.5** (Poole Theorem 3.19). *Let  $\mathbf{v}_1, \dots, \mathbf{v}_k \in \mathbb{R}^n$ . Then  $\text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_k)$  is a subspace of  $\mathbb{R}^n$ .*

*Proof.* For simplicity, let  $W = \text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_k)$ .

1. Since  $\mathbf{0} = 0\mathbf{v}_1 + \dots + 0\mathbf{v}_k$ , then  $\mathbf{0} \in W$ .
2. Let  $\mathbf{u} = c_1\mathbf{v}_1 + \dots + c_k\mathbf{v}_k$  and  $\mathbf{w} = d_1\mathbf{v}_1 + \dots + d_k\mathbf{v}_k$  be vectors in  $W$ . Then

$$\mathbf{u} + \mathbf{w} = (c_1 + d_1)\mathbf{v}_1 + \dots + (c_k + d_k)\mathbf{v}_k.$$

Since  $\mathbf{u} + \mathbf{w}$  is a linear combination of the vectors  $\mathbf{v}_i$ , then  $\mathbf{u} + \mathbf{w} \in W$ .

3. Let  $\mathbf{u}$  be as above and  $c \in \mathbb{R}$  be some scalar. Then

$$c\mathbf{u} = (cc_1)\mathbf{v}_1 + \dots + (cc_k)\mathbf{v}_k.$$

Since  $c\mathbf{u}$  is a linear combination of the vectors  $\mathbf{v}_i$ , then  $c\mathbf{u} \in W$ .

□

### 3.4.1 Subspaces Associated with Matrices

With the notion of a "subspace" in mind, let's try to revisit some ideas involving matrices. First, a new definition

**Definition.** Let  $A$  be an  $m \times n$  matrix.

1. The *column space* of  $A$  is a subspace  $\mathbb{R}^n$  spanned by the columns of  $A$ . We denote it as  $\text{Col}(A)$ .
2. The *row space* of  $A$  is a subspace of  $\mathbb{R}^m$  spanned by the rows of  $A$ . We denote it as  $\text{Row}(A)$

*Remark.* Since we will prefer to think about column vectors whenever possible, it may be more useful to define  $\text{Row}(A) := \text{Col}(A^T)$ .

**Theorem 3.4.6** (Poole Theorem 3.21). *Let  $A$  be an  $m \times n$  matrix and let  $N$  be the set of solutions to the homogeneous system  $A\mathbf{x} = \mathbf{0}$ . Then  $N$  is a subspace of  $\mathbb{R}^n$ .*

*Proof.* 1. Clear  $\mathbf{x} = \mathbf{0}$  is a solution to the homogeneous system.

2. Let  $\mathbf{x}_1, \mathbf{x}_2$  be in  $N$ . Then

$$A(\mathbf{x}_1 + \mathbf{x}_2) = A\mathbf{x}_1 + A\mathbf{x}_2 = \mathbf{0} + \mathbf{0} = \mathbf{0}.$$

so  $N$  is closed under addition.

3. Let  $\mathbf{x}$  be in  $N$  and  $k$  be a scalar. Then

$$A(k\mathbf{x}) = k(A\mathbf{x}) = k\mathbf{0} = \mathbf{0}$$

so  $N$  is closed under scalar multiplication. □

**Definition.**  $N$ , as above, is called the *null space* of  $A$ , and is denoted  $\text{Null}(A)$ . (In some texts, it is called the *kernel* of  $A$  and is denoted  $\ker(A)$ .)

**Example 3.4.7.** Compute  $\text{Col}(A)$ ,  $\text{Row}(A)$ , and  $\text{Null}(A)$  for  $A = \begin{bmatrix} 1 & 2 & 4 \\ 1 & 2 & 4 \\ 1 & 2 & 4 \end{bmatrix}$ .

Letting  $\mathbf{a}_i$  denote the  $i^{\text{th}}$  column of  $A$ , we see that  $\mathbf{a}_3 = 2\mathbf{a}_2 = 4\mathbf{a}_1$ , hence

$$\text{Col}(A) = \text{Span} \left( \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right).$$

Similarly, letting  $\mathbf{A}_i$  denote the  $i^{\text{th}}$  row of  $A$ , we see that  $\mathbf{A}_3 = \mathbf{A}_2 = \mathbf{A}_1$ , hence

$$\text{Row}(A) = \text{Span}([1, 2, 4]).$$

Examining the homogeneous system  $A\mathbf{x} = \mathbf{0}$ ,

$$[A \mid \mathbf{0}] = \left[ \begin{array}{ccc|c} 1 & 2 & 4 & 0 \\ 1 & 2 & 4 & 0 \\ 1 & 2 & 4 & 0 \end{array} \right] \xrightarrow[\substack{R_2 - R_1 \rightarrow R_2 \\ R_3 - R_1 \rightarrow R_3}]{R_3 - R_1 \rightarrow R_3} \left[ \begin{array}{ccc|c} 1 & 2 & 4 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

we see that  $A$  has rank 1 (hey wait, that's how many vectors span both  $\text{Row}(A)$  and  $\text{Col}(A)$ ... weird), hence there are two free variables in this system,  $x_2 = s$  and  $x_3 = t$ . We thus get that the solution set is of the form

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} -2s - 4t \\ s \\ t \end{bmatrix} = s \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} + t \begin{bmatrix} -4 \\ 0 \\ 1 \end{bmatrix}$$

hence

$$\text{Null}(A) = \text{Span} \left( \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} -4 \\ 0 \\ 1 \end{bmatrix} \right).$$

**Definition.** Let  $W$  be a subspace of a vector space and  $\mathcal{B} = \{w_1, \dots, w_k\}$  a set of vectors in  $W$ .  $\mathcal{B}$  is a *basis* for  $W$  if

1.  $W = \text{Span}(\mathcal{B})$  and
2.  $\mathcal{B}$  is a linearly independent set.

*Remark.* Since every vector space is a subspace of itself, this definition is valid for all vector spaces. We've merely stated it in terms of subspaces to make it clear that the basis vectors must each be contained in that subspace.



**Example 3.4.8.** The standard basis vectors  $\mathbf{e}_i$  in  $\mathbb{R}^n$  form a basis for  $\mathbb{R}^n$ . We refer to  $\mathcal{E} = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$  as the *standard basis* for  $\mathbb{R}^n$ .

**Example 3.4.9.** Find basis for the column space  $A = \begin{bmatrix} 1 & 1 & 3 & 1 & 6 \\ 2 & -1 & 0 & 1 & -1 \\ -3 & 2 & 1 & -2 & 1 \\ 4 & 1 & 6 & 1 & 3 \end{bmatrix}$ .

Notice that

$$\text{RREF}(A) = \begin{bmatrix} 1 & 0 & 1 & 0 & -1 \\ 0 & 1 & 2 & 0 & 3 \\ 0 & 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

Let  $\mathbf{a}_i$  represent the  $i^{\text{th}}$  column vector for  $A = [\mathbf{a}_1 \ \dots \ \mathbf{a}_5]$ . What the reduced row echelon form suggests is that

$$\mathbf{a}_3 = 1\mathbf{a}_1 + 2\mathbf{a}_2 \quad \text{and} \quad \mathbf{a}_5 = -1\mathbf{a}_1 + 3\mathbf{a}_2 + 4\mathbf{a}_4$$

(and you can check that this is true). Hence  $\text{Col}(A) = \text{Span}(\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_4)$ . Moreover,

$$\text{RREF}([\mathbf{a}_1 \ \mathbf{a}_2 \ \mathbf{a}_4 \mid \mathbf{0}]) = \left[ \begin{array}{ccc|c} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

and so  $\{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_4\}$  is a linearly independent set. Hence

$$\{\mathbf{a}_1, \mathbf{a}_2, \mathbf{a}_4\} = \left\{ \begin{bmatrix} 1 \\ 2 \\ -3 \\ 4 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \\ 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ -2 \\ 1 \end{bmatrix} \right\}$$

is a basis for  $\text{Col}(A)$ .

**Strategy for finding a basis for the column space of a matrix:**

1. Row reduce the matrix (just row-echelon form is fine)
2. Take as a basis every column (in the original matrix) which contains a leading 1/pivot.

**Example 3.4.10.** Find basis for the null space of  $A = \begin{bmatrix} 1 & 1 & 3 & 1 & 6 \\ 2 & -1 & 0 & 1 & -1 \\ -3 & 2 & 1 & -2 & 1 \\ 4 & 1 & 6 & 1 & 3 \end{bmatrix}$ .

Notice that

$$\text{RREF}(A) = \begin{bmatrix} 1 & 0 & 1 & 0 & -1 \\ 0 & 1 & 2 & 0 & 3 \\ 0 & 0 & 0 & 1 & 4 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

When we go to solve the system

$$A\mathbf{x} = A \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \mathbf{0}$$

we see that  $x_3$  and  $x_5$  are free variables and

$$\begin{aligned} x_1 &= -x_3 + x_5, \\ x_2 &= -2x_3 - 3x_5, \\ x_4 &= -4x_5. \end{aligned}$$

By setting  $x_3 = s$  and  $x_5 = t$ , we can parameterize the solution set as

$$\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \end{bmatrix} = \begin{bmatrix} -s + t \\ -2s - 3t \\ s \\ -4t \\ t \end{bmatrix} = s \begin{bmatrix} -1 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix} + t \begin{bmatrix} 1 \\ -3 \\ 0 \\ -4 \\ 1 \end{bmatrix} = s\mathbf{v}_1 + t\mathbf{v}_2.$$

Clearly  $\text{Null}(A) = \text{Span}(\mathbf{v}_1, \mathbf{v}_2)$  and it is straightforward to check that  $\{\mathbf{v}_1, \mathbf{v}_2\}$  are linearly independent, hence

$$\{\mathbf{v}_1, \mathbf{v}_2\} = \left\{ \begin{bmatrix} -1 \\ -2 \\ 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ -3 \\ 0 \\ -4 \\ 1 \end{bmatrix} \right\}$$

is a basis for  $\text{Null}(A)$ .

### Strategy for finding a basis for the null space of a matrix $A$ :

1. Row reduce the matrix.
2. Solve the system  $A\mathbf{x} = \mathbf{0}$ .
3. Solve for all columns with leading 1's/pivots in terms of free variables.
4. Write the solution set as a parameterized linear combination of  $k$  vectors (where  $k$  is the number of free variables).
5. Take these  $k$  vectors as a basis for  $\text{Null}(A)$ .

### How many bases can a vector space have?

Infinitely-many:

**Exercise 3.4.1.** Show that, for any nonzero real numbers  $m, n$ , the set  $\left\{ \begin{bmatrix} m \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ n \end{bmatrix} \right\}$  is a basis for  $\mathbb{R}^2$ .

Maybe the better question is

**How many vectors must a basis have? Can two different bases have different numbers of vectors?**

**Example 3.4.11.** Suppose  $V$  is a subspace of  $\mathbb{R}^3$  and has two different bases,  $\mathcal{B}_1 = \{\mathbf{u}_1, \mathbf{u}_2\}$  and  $\mathcal{B}_2 = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$ .

Since  $\mathcal{B}_1$  is a basis, we can write each of the  $\mathcal{B}_2$ -basis vectors as a linear combination of the  $\mathcal{B}_1$ -basis vectors

$$\mathbf{v}_1 = a_{11}\mathbf{u}_1 + a_{21}\mathbf{u}_2$$

$$\mathbf{v}_2 = a_{12}\mathbf{u}_1 + a_{22}\mathbf{u}_2$$

$$\mathbf{v}_3 = a_{13}\mathbf{u}_1 + a_{23}\mathbf{u}_2$$

Now if we consider the vector equation

$$\mathbf{0} = x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + x_3\mathbf{v}_3$$

it *should be* that the only solution is when each of the  $x_i = 0$  (since the we claim the  $\mathbf{v}_i$ 's are linearly independent). Notice, however, that

$$\mathbf{0} = x_1\mathbf{v}_1 + x_2\mathbf{v}_2 + x_3\mathbf{v}_3$$

$$\mathbf{0} = x_1(a_{11}\mathbf{u}_1 + a_{21}\mathbf{u}_2) + x_2(a_{12}\mathbf{u}_1 + a_{22}\mathbf{u}_2) + x_3(a_{13}\mathbf{u}_1 + a_{23}\mathbf{u}_2)$$

$$\mathbf{0} = (a_{11}x_1 + a_{12}x_2 + a_{13}x_3)\mathbf{u}_1 + (a_{21}x_1 + a_{22}x_2 + a_{23}x_3)\mathbf{u}_2$$

Since the  $\mathbf{u}_i$ 's are linearly independent, we must have that

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + a_{13}x_3 = 0 \\ a_{21}x_1 + a_{22}x_2 + a_{23}x_3 = 0 \end{cases}$$

which is a homogeneous linear system in 3 variables (and only 2 equations), hence there are infinitely many solutions, and hence  $\mathcal{B}_2$  is a linearly dependent set and not a basis.

### 3.4.2 Dimension and Rank

**Theorem 3.4.12** (Poole Theorem 3.23 - The Basis Theorem). *Let  $V$  be a vector space with two different bases  $\mathcal{B}_1$  and  $\mathcal{B}_2$ . Then  $\mathcal{B}_1$  and  $\mathcal{B}_2$  have the same number of vectors.*

Because the number of vectors in the basis is invariant of the choice of basis, we can define the following term.

**Definition.** The *dimension* of a vector space  $V$  is the number of vectors in a basis for  $V$ . We denote this  $\dim(V)$ .

*Remark.* The trivial vector space  $\{\mathbf{0}\}$  is defined to have dimension 0.

**Example 3.4.13.**  $\dim(\mathbb{R}^n) = n$ .

**Example 3.4.14.** Let  $V = \text{Span} \left( \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix} \right)$  be a subspace of  $\mathbb{R}^3$ . Clearly the first vector is a linear combination of the other two (which are linearly independent), so  $\dim(V) = 2$ .

**Theorem 3.4.15** (Poole Theorem 3.24). *For a matrix  $A$ ,  $\dim(\text{Row}(A)) = \dim(\text{Col}(A))$ .*

**Definition.** The *rank* of a matrix  $A$  is the dimension of its column space (denoted  $\text{Rank}(A)$ ). If  $A$  has size  $n \times n$  and  $\text{Rank}(A) = n$ , then sometimes we say that  $A$  has *full rank*.

*Remark.* This new notion of rank still agrees with our old notion, because the number of linearly independent rows in  $A$  is the same as the number of nonzero rows in  $RREF(A)$  and  $\dim(\text{Row}(A)) = \dim(\text{Col}(A))$ .

**Theorem 3.4.16** (Poole Theorem 3.25). For any matrix  $A$ ,  $\text{Rank}(A) = \text{Rank}(A^T)$

**Definition.** The *nullity* of a matrix  $A$  is the dimension of its null space. We denote it by  $\text{nullity}(A)$ .

**Theorem 3.4.17** (Poole Theorem 3.26 - Rank-Nullity). If  $A$  is an  $m \times n$  matrix, then  $\text{Rank}(A) + \text{nullity}(A) = n$ .

**Example 3.4.18.** Show that  $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \\ -2 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ -3 \end{bmatrix} \right\}$  is a basis for  $\mathbb{R}^3$

We consider the matrix  $A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 1 \\ 1 & -2 & -3 \end{bmatrix}$  and compute its rank.

$$\begin{aligned} \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 1 \\ 1 & -2 & -3 \end{bmatrix} &\xrightarrow{R_3 - R_1 \rightarrow R_3} \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 1 \\ 0 & -4 & -6 \end{bmatrix} \\ &\xrightarrow{R_3 + 4R_2 \rightarrow R_3} \begin{bmatrix} 1 & 2 & 3 \\ 0 & 1 & 1 \\ 0 & 0 & -2 \end{bmatrix} \end{aligned}$$

Thus  $\text{Rank}(A) = 3$ . Hence the columns of  $A$  span a 3-dimensional subspace of  $\mathbb{R}^3$ , i.e.  $\text{Col}(A) = \mathbb{R}^3$ . Since the columns of  $A$  are linearly independent and span  $\mathbb{R}^3$ ,  $\mathcal{B}$  is a basis for  $\mathbb{R}^3$ .

**Example 3.4.19.** Show that  $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 3 \\ -7 \\ 3 \end{bmatrix} \right\}$  is not a basis for  $\mathbb{R}^3$ .

Employ the same technique as in the previous example, see that the rank is only 2.

**Theorem 3.4.20** (Fundamental Theorem of Invertible Matrices, Pt II). Suppose  $A$  is an  $n \times n$  matrix. The following are equivalent:

- a.  $A$  is invertible.
- ⋮
- g. The column vectors of  $A$  span  $\mathbb{R}^n$ .
- h. The column vectors of  $A$  form a basis for  $\mathbb{R}^n$ .
- i. The row vectors of  $A$  are linearly independent.
- j. The row vectors of  $A$  span  $\mathbb{R}^n$ .
- k. The row vectors of  $A$  form a basis for  $\mathbb{R}^n$ .
- l.  $\text{Rank}(A) = n$
- m.  $\text{nullity}(A) = 0$

### 3.4.3 Coordinates

**Theorem 3.4.21.** Let  $V$  be a vector space with an ordered basis  $\mathcal{B} = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ . For every vector  $\mathbf{u} \in V$ , there is a unique linear combination of  $\mathcal{B}$ -basis vectors such that

$$\mathbf{u} = c_1\mathbf{v}_1 + \dots + c_k\mathbf{v}_k.$$

**Definition.** The  $c_i$  in the previous theorem are called the *coordinates of  $\mathbf{u}$  with respect to  $\mathcal{B}$*  and the column vector

$$[\mathbf{u}]_{\mathcal{B}} = \begin{bmatrix} c_1 \\ \vdots \\ c_k \end{bmatrix}$$

is called the *coordinate vector of  $\mathbf{u}$  with respect to  $\mathcal{B}$* .

**Example 3.4.22.** Let  $P(0,0)$  and  $Q(3,1)$  be points in the plane and consider the vector  $\mathbf{v} = \overrightarrow{PQ}$ . Given the standard basis  $\mathcal{E} = \{\mathbf{e}_1, \mathbf{e}_2\}$  for  $\mathbb{R}^2$ , we can write

$$\mathbf{v} = 3\mathbf{e}_1 + 1\mathbf{e}_2$$

hence

$$[\mathbf{v}]_{\mathcal{E}} = \begin{bmatrix} 3 \\ 1 \end{bmatrix}.$$


**Example 3.4.23.** With  $P, Q, \mathbf{v}$  as above, we consider now the basis  $\mathcal{B} = \{\mathbf{b}_1, \mathbf{b}_2\} = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$  for  $\mathbb{R}^2$ . Since

$$\mathbf{v} = 2\mathbf{b}_1 + 1\mathbf{b}_2$$


then

$$[\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}.$$

Visually, the previous two examples just give a bit of formality to the hand-wavy "coordinate grid" discussion from Section 1.1



$\mathbf{v}$  in the  $\mathcal{E}$ -basis.



$\mathbf{v}$  in the  $\mathcal{B}$ -basis.

*Remark.* We typically don't write the subscript  $\mathcal{E}$  for vectors when they are written in the standard basis.

### Section 3.4 Exercises

1. Let  $V = \text{Span} \left( \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 2 \\ -1 \\ 0 \end{bmatrix} \right)$  and define  $W$  to be the set of all vectors  $\mathbf{x} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix}$  in  $\mathbb{R}^3$  satisfying
- $$x_1 + 2x_2 + 3x_3 = 0 \quad \text{and} \quad 2x_1 - x_2 = 0.$$

$W$  is known as the **orthogonal complement** of  $V$  in  $\mathbb{R}^3$ .

- Show that  $W$  is a subspace of  $\mathbb{R}^3$ .
  - Find a vector  $\mathbf{w}$  so that  $W = \text{Span}(\mathbf{w})$ .
  - Find a matrix  $A$  so that  $W = \text{Null}(A)$ .
  - With the matrix  $A$  you found in part (c), find a basis for  $\text{Col}(A^T)$ .
2. Suppose  $A$  is a  $5 \times 7$  matrix and  $\text{Rank}(A) = 3$ .
- $\text{Null}(A)$  is a subspace of  $\mathbb{R}^k$  for some  $k$ . What is the value of  $k$ ?
  - What is the dimension of  $\text{Null}(A)$ ?
3. Recall from Section 1.1 that  $\mathcal{P}^n$ , the set of all polynomials of degree at most  $n$  (with indeterminate  $x$  and real coefficients), is a real vector space, where “vector addition” is just the usual polynomial addition, and “scalar multiplication” is the usual scalar multiplication:

$$\begin{aligned} \mathbf{u} + \mathbf{v} &= (u_0 + v_0) + (u_1 + v_1)x + \cdots + (u_n + v_n)x^n \\ k\mathbf{v} &= kv_0 + kv_1x + \cdots + kv_nx^n \end{aligned}$$

- Show that  $\mathcal{P}^2$  is a subspace of  $\mathcal{P}^3$ .
- Find a basis for  $\mathcal{P}^2$  and provide justification that it is indeed a basis. (Since each “vector” in  $\mathcal{P}^2$  is a polynomial, so too will be the basis “vectors”.)

### 6.3 Change of Basis

Let  $\mathcal{E} = \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$  be the standard basis for  $\mathbb{R}^2$  and let  $\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$  another basis for  $\mathbb{R}^2$ . We saw that a particular vector  $\mathbf{v} \in \mathbb{R}^2$  had coordinate representation

$$[\mathbf{v}]_{\mathcal{E}} = \begin{bmatrix} 3 \\ 1 \end{bmatrix} \quad \text{and} \quad [\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}.$$

Notice that

$$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} [\mathbf{v}]_{\mathcal{B}} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \end{bmatrix} = [\mathbf{v}]_{\mathcal{E}}.$$

Multiplying by a matrix had the effect of changing the basis in which  $\mathbf{v}$  was written!

**Definition.** Let  $\mathcal{B} = \{\mathbf{b}_1, \dots, \mathbf{b}_n\}$  and  $\mathcal{C} = \{\mathbf{c}_1, \dots, \mathbf{c}_n\}$  be two bases for  $\mathbb{R}^n$ . The  $n \times n$  matrix

$$P_{\mathcal{C} \leftarrow \mathcal{B}} = \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & \cdots & [\mathbf{b}_n]_{\mathcal{C}} \end{bmatrix}.$$

is called the *change-of-basis matrix* from  $\mathcal{B}$  to  $\mathcal{C}$ . It has the effect

$$P_{\mathcal{C} \leftarrow \mathcal{B}} [\mathbf{v}]_{\mathcal{B}} = [\mathbf{v}]_{\mathcal{C}}.$$

*Fact.* Given two different bases for  $\mathbb{R}^n$ ,  $\mathcal{B}$  and  $\mathcal{C}$ , the following are true

- $(P_{\mathcal{C} \leftarrow \mathcal{B}})^{-1} = P_{\mathcal{B} \leftarrow \mathcal{C}}$ , and
- $P_{\mathcal{C} \leftarrow \mathcal{E}} P_{\mathcal{E} \leftarrow \mathcal{B}} = P_{\mathcal{C} \leftarrow \mathcal{B}}$ ,

where  $\mathcal{E}$  is the standard basis.

**Example 6.3.1.** Let  $\mathcal{B} = \left\{ \mathbf{b}_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \mathbf{b}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}$  and  $\mathcal{C} = \left\{ \mathbf{c}_1 = \begin{bmatrix} 1 \\ -2 \end{bmatrix}, \mathbf{c}_2 = \begin{bmatrix} -3 \\ 3 \end{bmatrix} \right\}$  be bases for  $\mathbb{R}^2$ . Compute the change of basis matrix  $P_{\mathcal{C} \leftarrow \mathcal{B}}$ .

We need to find  $[\mathbf{b}_1]_{\mathcal{C}}$  and  $[\mathbf{b}_2]_{\mathcal{C}}$ .

$$\begin{aligned} \mathbf{b}_1 = -2\mathbf{c}_2 - \mathbf{c}_1 &\implies [\mathbf{b}_1]_{\mathcal{C}} = \begin{bmatrix} -2 \\ -1 \end{bmatrix} \\ \mathbf{b}_2 = 0\mathbf{c}_2 - \frac{1}{3}\mathbf{c}_2 &\implies [\mathbf{b}_2]_{\mathcal{C}} = \begin{bmatrix} 0 \\ -\frac{1}{3} \end{bmatrix} \end{aligned}$$

so

$$P_{\mathcal{C} \leftarrow \mathcal{B}} = \begin{bmatrix} [\mathbf{b}_1]_{\mathcal{C}} & [\mathbf{b}_2]_{\mathcal{C}} \end{bmatrix} = \begin{bmatrix} -2 & 0 \\ -1 & -\frac{1}{3} \end{bmatrix}.$$



## 3.6 Introduction to Linear Transformations

**Definition.** A *transformation* (aka *function* or *map*) is a function  $T$  with domain  $\mathbb{R}^n$  and codomain  $\mathbb{R}^m$ , written

$$T : \mathbb{R}^n \rightarrow \mathbb{R}^m.$$

$T$  is a *linear transformation* if

1.  $T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$  for all  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^m$ , and
2.  $T(k\mathbf{v}) = kT(\mathbf{v})$  for all scalars  $k \in \mathbb{R}$  and vectors  $\mathbf{v} \in \mathbb{R}^m$ .

**Example 3.6.1** (Identity transformation).  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  given by  $T(\mathbf{v}) = \mathbf{v}$  is a linear transformation.

**Example 3.6.2** (Trivial transformation).  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  given by  $T(\mathbf{v}) = \mathbf{0}$  is a linear transformation.

**Example 3.6.3.** Consider  $T : \mathbb{R} \rightarrow \mathbb{R}$  given by  $T(x) = mx + b$ , where  $b$  is some constant. Then

1.  $T(x_1 + x_2) = mx_1 + mx_2 + b = (mx_1 + b) + (mx_2 + b) = T(x_1) + T(x_2)$  (precisely when  $b = 0$ ), and
2.  $T(kx) = mkx + b = k(mx + b) = kT(x)$  (precisely when  $b = 0$ )

so  $T$  is a linear transformation only when  $b = 0$ .

**Example 3.6.4.** Consider the map

$$\begin{aligned} T_A : \mathbb{R}^2 &\rightarrow \mathbb{R}^3 \\ T_A(\mathbf{x}) &= A\mathbf{x} \end{aligned}$$

where  $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \\ 5 & 6 \end{bmatrix}$ . It follows from matrix arithmetic rules that This is a linear transformation because one can easily check that

1.  $T_A(\mathbf{x}_1 + \mathbf{x}_2) = A(\mathbf{x}_1 + \mathbf{x}_2) = A\mathbf{x}_1 + A\mathbf{x}_2 = T_A(\mathbf{x}_1) + T_A(\mathbf{x}_2)$
2.  $T_A(k\mathbf{x}) = A(k\mathbf{x}) = kA\mathbf{x} = kT_A(\mathbf{x})$

The above argument actually shows the following more general result

**Theorem 3.6.5.** If  $A \in \mathbb{R}^{m \times n}$ , then the transformation

$$\begin{aligned} T_A : \mathbb{R}^n &\rightarrow \mathbb{R}^m \\ T_A(\mathbf{x}) &= A\mathbf{x} \end{aligned}$$

is a linear transformation.

Notice that, explicitly, we could write

$$T_A\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} x + 2y \\ 3x + 4y \\ 5x + 6y \end{bmatrix} = x \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} + y \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix}.$$

The range of  $T_A$  then is just

$$\begin{aligned} \text{Range}(T_A) &= \left\{ T_A \left( \begin{bmatrix} x \\ y \end{bmatrix} \right) : \text{where } x, y \in \mathbb{R} \right\} \\ &= \left\{ x \begin{bmatrix} 1 \\ 3 \\ 5 \end{bmatrix} + y \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix} : \text{where } x, y \in \mathbb{R} \right\} = \text{Col}(A) \end{aligned}$$

As it turns out, we can write every linear transformation as multiplication by a matrix.


**Theorem 3.6.6.** *Let  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  be a linear transformation. Then we can write  $T(\mathbf{x}) = A\mathbf{x}$  where  $A$  is the  $m \times n$  matrix whose  $i^{\text{th}}$  column is the column vector  $T(\mathbf{e}_i)$ , i.e.*

$$A = \left[ \begin{array}{c|c|c} T(\mathbf{e}_1) & \cdots & T(\mathbf{e}_m) \end{array} \right].$$

**Definition.** The matrix in the above theorem is called the *standard matrix of  $T$* . We may sometimes write  $[T]$  to denote the standard matrix of  $T$ .

### 3.6.1 Types of Linear Transformations of $\mathbb{R}^2$

**Example 3.6.7.** A Reflection about the  $x$ -axis,  $R_x$ , is a linear transformation of  $\mathbb{R}^2$ .



Explicitly, it sends a points  $(x, y)$  to  $(x, -y)$ , hence the transformation is given by


$$R_x \left( \begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} x \\ -y \end{bmatrix}$$

The standard matrix for  $R_x$  is thusly given by

$$[R_x] = \left[ R_x \left( \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right) \quad R_x \left( \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right) \right] = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$$

**Exercise 3.6.1.** Show that reflection about the  $y$ -axis is also a linear transformation and find the standard matrix for this.

**Example 3.6.8.** Rotation by an angle  $\theta$  about the origin,  $R_\theta$ , is a linear transformation of  $\mathbb{R}^2$ .




For this one, we'll first find the standard transformation matrix. Note that  $\mathbf{e}_1 = [\cos(0), \sin(0)]^T$  and  $\mathbf{e}_2 = [\cos(\frac{\pi}{2}), \sin(\frac{\pi}{2})]^T$ . So rotation by an angle  $\theta$  should add  $\theta$  to the angle arguments of sine and cosine, i.e.

$$\begin{aligned} [R_\theta] &= \left[ R_\theta \left( \begin{bmatrix} \cos(0) \\ \sin(0) \end{bmatrix} \right) \quad R_\theta \left( \begin{bmatrix} \cos(\frac{\pi}{2}) \\ \sin(\frac{\pi}{2}) \end{bmatrix} \right) \right] \\ &= \begin{bmatrix} \cos(0 + \theta) & \cos(\frac{\pi}{2} + \theta) \\ \sin(0 + \theta) & \sin(\frac{\pi}{2} + \theta) \end{bmatrix} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix} \end{aligned}$$

hence the linear transformation is given by

$$R_\theta \left( \begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} x \cos(\theta) - y \sin(\theta) \\ x \sin(\theta) + y \cos(\theta) \end{bmatrix}$$

**Example 3.6.9.** A **dilation** (with **dilation factor**  $k$ ) is a transformation  $D_k$  that expands out from the origin by a factor of  $k$ .




Explicitly, it sends a point  $(x, y)$  to a point  $(kx, ky)$  so for vectors,

$$D_k \left( \begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} kx \\ ky \end{bmatrix}$$

The standard matrix for  $D_k$  is given by

$$[D_k] = \left[ D_k \left( \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right) \quad D_k \left( \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right) \right] = \begin{bmatrix} k & 0 \\ 0 & k \end{bmatrix}.$$

**Example 3.6.10.** A **horizontal shear** (with **shear factor**  $m$ ),  $S_m$ , is a transformation that slides the top edge of the unit square  $m$  units to the right (making a parallelogram).



In particular, it sends  $(x, y)$  to the point  $(x + my, y)$ ,

$$S_m \left( \begin{bmatrix} x \\ y \end{bmatrix} \right) = \begin{bmatrix} x + my \\ y \end{bmatrix}$$

The standard matrix for  $S_m$  is given by

$$[S_m] = \left[ S_m \left( \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right) \quad S_m \left( \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right) \right] = \begin{bmatrix} 1 & m \\ 0 & 1 \end{bmatrix}$$

**Theorem 3.6.11.** If  $T_A : \mathbb{R}^m \rightarrow \mathbb{R}^n$  and  $T_B : \mathbb{R}^n \rightarrow \mathbb{R}^p$  are two linear transformations (with standard matrices  $A$  and  $B$ ), then

$$T_B \circ T_A = T_{BA}.$$

**Definition.** A transformation  $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is *one-to-one* (or *injective*) if, for every vector  $\mathbf{v} \in \mathbb{R}^n$ , there is a unique  $\mathbf{w} \in \mathbb{R}^m$  for which  $T(\mathbf{v}) = \mathbf{w}$ .  $T$  is *onto* (or *surjective*) if, for every  $\mathbf{w} \in \mathbb{R}^m$ , there is at least one  $\mathbf{v} \in \mathbb{R}^n$  for which  $T(\mathbf{v}) = \mathbf{w}$ . The *kernel* of  $T$  is the collection of vectors  $\mathbf{x}$  in  $\mathbb{R}^n$  for which  $T(\mathbf{x}) = \mathbf{0}$ .

*Remark.* Maybe the right way to think about it is this: “one-to-one” means that the range of  $T$  is a copy of  $\mathbb{R}^n$  living inside of  $\mathbb{R}^m$ , and “onto” means that the range of  $T$  is all of  $\mathbb{R}^m$ .

**Theorem 3.6.12.** Let  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^m$ .

1.  $T_A$  is one-to-one if and only if  $\text{Rank}(A) = n$ .
2.  $T_A$  is onto if and only if  $\text{Rank}(A) = m$ .
3.  $\ker(T_A) = \text{Null}(A)$ .

**Definition.** Two transformations  $S, T : \mathbb{R}^n \rightarrow \mathbb{R}^m$  are *inverses* if, for every  $\mathbf{v} \in \mathbb{R}^n$ ,

$$S \circ T(\mathbf{v}) = T \circ S(\mathbf{v}) = \mathbf{v}$$

**Theorem 3.6.13.** If  $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$  has an inverse, then the standard matrix  $[T]$  is invertible and  $[T^{-1}] = [T]^{-1}$


**Theorem 3.6.14.**  $A$  is invertible if and only if  $T_A : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is both one-to-one and onto.

## 3.7 Applications

### 3.7.1 Markov Chains

A Markov chain is just a process by which one models probabilistic scenarios

**Example 3.7.1.** Researchers have found that Democrat (D) voters are 70% likely to continue voting Democrat in the next election, 10% likely to vote Republican in the next election, and 20% likely to vote Independent in the next election. Similar data was compiled for Republican (R) and Independent (I) voters, and can be modeled in the following graph:



If there are  $D_0$  Democrat voters,  $R_0$  Republican voters, and  $I_0$  Independent voters in this current election cycle, how many of each will there be for the next election cycle? We can write

$$\begin{aligned}D_1 &= 0.7(D_0) + 0.1(R_0) + 0.3(I_0) \\R_1 &= 0.1(D_0) + 0.8(R_0) + 0.2(I_0) \\I_1 &= 0.2(D_0) + 0.1(R_0) + 0.5(I_0)\end{aligned}$$

or, as a matrix/vectors

$$\mathbf{x}_1 = \begin{bmatrix} D_1 \\ R_1 \\ I_1 \end{bmatrix} = \begin{bmatrix} 0.7 & 0.1 & 0.3 \\ 0.1 & 0.8 & 0.2 \\ 0.2 & 0.1 & 0.5 \end{bmatrix} \begin{bmatrix} D_0 \\ R_0 \\ I_0 \end{bmatrix} = A\mathbf{x}_0$$

It follows that, after  $k$  elections cycles,  $\mathbf{x}_k = A^k\mathbf{x}_0$ .

**Definition.** The vector  $\mathbf{x}_0$  is known as the *initial state vector*,  $A$  is known as the *transition matrix*, and the entire process above is called a *Markov chain* (with 3 states). If the transition matrix has columns with positive entries that all sum to 1, it is called a *stochastic matrix*, the columns are called *probability vectors*, and the  $(i, j)$ -entry represents the probability that object in state  $i$  (e.g. voting preference) switches to state  $j$ .

Below are a few computations for different  $k$ -values, assuming  $D_0 = 1000$ ,  $R_0 = 800$  and  $I_0 = 300$ :

$$\mathbf{x}_1 = \begin{bmatrix} 870 \\ 800 \\ 430 \end{bmatrix}, \quad \mathbf{x}_{20} \approx \begin{bmatrix} 764 \\ 859 \\ 477 \end{bmatrix}, \quad \mathbf{x}_{100} \approx \begin{bmatrix} 764 \\ 859 \\ 477 \end{bmatrix}, \quad \mathbf{x}_{1000} \approx \begin{bmatrix} 764 \\ 859 \\ 477 \end{bmatrix}$$

What we notice is that the vector  $\mathbf{x}_k = A^k \mathbf{x}_0$  seems to stop changing as  $k \rightarrow \infty$ .

If you're interested in playing around with this yourself, say with different initial conditions or a different number of steps in the Markov chain process, you can use the MATLAB code below:

```
transMat = [ 0.7 0.3 0.1 ; 0.1 0.2 0.8 ; 0.2 0.5 0.1 ]; %transition matrix
x0 = [ 1000 ; 800 ; 300 ]; %initial state vector [D0,R0,I0]
maxLoop = 100; %number of iterations in Markov chain

for k=1:maxLoop
    transpose(transMat^k*x0) %outputs xk = [Dk,Rk,Ik], kth step in Markov chain process
end
```

What's fascinating is that this limit doesn't change as long as  $D_0 + R_0 + I_0 = 2100$  (really, try it).

**Definition.** In a Markov chain process with transition matrix  $A$ , a vector  $\mathbf{x}$  for which  $A\mathbf{x} = \mathbf{x}$  is called a *steady state vector*.

*Remark.* Every Markov chain has a steady state vector, but unless all of the entries of the transition matrix are positive, it may not be the limit  $\lim_{k \rightarrow \infty} A^k \mathbf{x}_0$  (such a limit may not even exist).

Even more interesting - in the last example, not only did the limit not change under mild assumptions about the initial state vector, it turns out that no matter what nonzero vector we choose for  $\mathbf{x}_0$ , it will always be the case that

$$\lim_{k \rightarrow \infty} A^k \mathbf{x}_0 \approx \lambda \begin{bmatrix} 764 \\ 859 \\ 477 \end{bmatrix}$$

(for some real number  $\lambda$ ).

## 4.1 Introduction to Eigenvalues and Eigenvectors

**Definition.** Let  $A$  be an  $n \times n$  matrix. A scalar  $\lambda$  is an *eigenvalue* of  $A$  if there is a nonzero vector  $\mathbf{v} \in \mathbb{R}^n$  so that  $A\mathbf{v} = \lambda\mathbf{v}$ . Such a vector is called an *eigenvector* of  $A$  corresponding to  $\lambda$ .

*Remark.* The prefix *eigen-* is not a name, but is derived from German and means “special” or “characteristic.”

**Example 4.1.1.** In the Markov chain example, the steady state vector was precisely the vector  $\mathbf{x}$  satisfying  $A\mathbf{x} = \mathbf{x}$ , so  $\mathbf{x}$  was an eigenvector of  $A$  corresponding to the eigenvalue 1.

**Example 4.1.2.**  $A = \begin{bmatrix} -6 & 3 \\ 4 & 5 \end{bmatrix}$  has eigenvector  $\begin{bmatrix} -3 \\ 1 \end{bmatrix}$ . Find the corresponding eigenvalue.

$$\begin{bmatrix} -6 & 3 \\ 4 & 5 \end{bmatrix} \begin{bmatrix} -3 \\ 1 \end{bmatrix} = \begin{bmatrix} 21 \\ -7 \end{bmatrix} = \lambda \begin{bmatrix} -3 \\ 1 \end{bmatrix}$$

hence the corresponding eigenvalue is  $\lambda = -7$ .

**Example 4.1.3.** Show that 6 is another eigenvalue of  $A = \begin{bmatrix} -6 & 3 \\ 4 & 5 \end{bmatrix}$  and find its corresponding eigenvector.

We need to find a vector  $\mathbf{v}$  such that

$$A\mathbf{v} = 6\mathbf{v} \quad \implies \quad A\mathbf{v} - 6\mathbf{v} = \mathbf{0} \quad \implies \quad (A - 6I)\mathbf{v} = \mathbf{0}$$

so really we need to compute  $\text{Null}(A - 6I)$ .

$$[A - 6I \mid \mathbf{0}] = \left[ \begin{array}{cc|c} -12 & 3 & 0 \\ 4 & -1 & 0 \end{array} \right] \xrightarrow{\text{RREF}} \left[ \begin{array}{cc|c} 4 & -1 & 0 \\ 0 & 0 & 0 \end{array} \right]$$

So a vector  $\mathbf{v}$  is in  $\text{Null}(A - 6I)$  if it has the form  $\begin{bmatrix} t \\ 4t \end{bmatrix}$ . As such, any nonzero vector of this form is an eigenvector of  $A$  corresponding to 6.

**Definition.** Let  $A$  be an  $n \times n$  matrix and  $\lambda$  an eigenvalue with corresponding eigenvectors  $\mathbf{v}_1, \dots, \mathbf{v}_k$ . The *eigenspace* corresponding to  $\lambda$  is

$$E_\lambda := \text{Span}(\mathbf{v}_1, \dots, \mathbf{v}_k).$$


*Remark.* It may at first seem surprising that any linear combination of the above  $\mathbf{v}_i$ 's is still an eigenvector for  $\lambda$ , but it is a straightforward computation to see that it is true:

$$\begin{aligned} A(c_1\mathbf{v}_1 + \dots + c_k\mathbf{v}_k) &= c_1A\mathbf{v}_1 + \dots + c_kA\mathbf{v}_k \\ &= c_1\lambda\mathbf{v}_1 + \dots + c_k\lambda\mathbf{v}_k \\ &= \lambda(c_1\mathbf{v}_1 + \dots + c_k\mathbf{v}_k). \end{aligned}$$


**Example 4.1.4.** In the previous example,  $E_6 = \text{Span}\left(\begin{bmatrix} 1 \\ 4 \end{bmatrix}\right)$

Geometrically, eigenvectors tell us about lines that are preserved by the linear transformation (since all that happens is a vector in that line is scaled).

**Example 4.1.5.** Using  $A$  as before, notice that the vectors  $[-3, 1]^T$  and  $[1, 4]^T$  do not change direction after a transformation, but are merely scaled.



Before applying transformation  $A$ .



After applying transformation  $A$ .

### How do I compute eigenvalues for a given matrix $A$ ?

**Definition.** If  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ , the *determinant* of  $A$  is

$$\det(A) = ad - bc$$

We'll define the determinants for all  $n \times n$  matrices in the next section.

An eigenvector for the eigenvalue  $\lambda$  is a vector in  $\text{Null}(A - \lambda I)$ . By the Fundamental Theorem of Invertible matrices tells us that  $\text{nullity}(A - \lambda I) \neq 0$  if and only if  $A - \lambda I$  is *not* invertible. Moreover, from a previous theorem, a  $2 \times 2$  matrix is *not* invertible if and only if  $\det(A - \lambda I) = 0$ . Since  $\det(A - \lambda I)$  is a polynomial in  $\lambda$ , this means that eigenvalues are precisely the zeroes of this polynomial.

**Example 4.1.6.** Find the eigenvalues for  $A = \begin{bmatrix} -6 & 3 \\ 4 & 5 \end{bmatrix}$ .

$$\begin{aligned} 0 &= \det(A - \lambda I) \\ &= \det\left(\begin{bmatrix} -6 - \lambda & 3 \\ 4 & 5 - \lambda \end{bmatrix}\right) \\ &= (-6 - \lambda)(5 - \lambda) - 12 \\ &= \lambda^2 + \lambda - 42. \end{aligned}$$

Using your favorite method of solving for the zeroes of this polynomial, we exactly see that its zeroes are  $\lambda = -6, 7$ , which are precisely the eigenvalues we expected to get from the previous examples.



**Example 4.1.7.** Find the eigenvalues and corresponding eigenspaces for the scalar matrix  $A = \begin{bmatrix} 7 & 0 \\ 0 & 7 \end{bmatrix}$ .

$$0 = \det(A - \lambda I) = \det\left(\begin{bmatrix} 7 - \lambda & 0 \\ 0 & 7 - \lambda \end{bmatrix}\right) = (7 - \lambda)(7 - \lambda).$$

This matrix has a single eigenvalue  $\lambda = 7$  and

$$E_7 = \text{Null}(A - 7I) = \text{Null}\left(\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}\right) = \text{Span}\left(\begin{bmatrix} 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}\right).$$

**Exercise 4.1.1.** Find the eigenvalues and corresponding eigenspaces for the horizontal shear matrix

$$A = \begin{bmatrix} 1 & 7 \\ 0 & 1 \end{bmatrix}.$$

## APPENDIX C Complex Numbers

Some matrices may have complex eigenvalues (what does that mean geometrically? hmmm...), so below is an list of important properties of complex numbers.

Let  $i = \sqrt{-1}$ , the so-called *imaginary unity*. A *complex number* is a number  $z = a + bi$  where  $a$  and  $b$  are real numbers.  $a$  is called the *real part* and  $b$  is called the *imaginary part* of  $z$ . The *conjugate of  $z$*  is the complex number  $\bar{z} = a - bi$ .

*Addition* of two complex numbers  $z_1$  and  $z_2$  is done via by adding the real and imaginary parts separately:

$$z_1 + z_2 = (a_1 + b_1i) + (a_2 + b_2i) = (a_1 + a_2) + i(b_1 + b_2).$$

*Multiplication* of complex numbers follows the usual distributive law:

$$\begin{aligned} z_1 z_2 &= (a_1 + b_1i)(a_2 + b_2i) \\ &= a_1 a_2 + a_1 b_2 i + a_2 b_1 i + b_1 b_2 i^2 \\ &= (a_1 a_2 - b_1 b_2) + (a_1 b_2 + a_2 b_1) i. \end{aligned}$$

Noting that  $z\bar{z}$  is a real number for any  $z$ , *division* of complex numbers is done by multiplying by the conjugate and scaling by  $1/z\bar{z}$ :

$$\frac{z_1}{z_2} = \frac{z_1 \bar{z}_2}{z_2 \bar{z}_2} = \frac{1}{z_2 \bar{z}_2} (z_1 \bar{z}_2) = \frac{1}{a_2^2 + b_2^2} ((a_1 a_2 + b_1 b_2) + (-a_1 b_2 + a_2 b_1) i)$$

**Exercise.** To each complex number  $z = a + bi$  we can associate the matrix  $[z] = \begin{bmatrix} a & -b \\ b & a \end{bmatrix}$ . Show that the product  $z_1 z_2$  can be derived from multiplying the associated matrices  $[z_1][z_2]$ .

### Matrices with complex entries

Although we won't see them in this course, matrices with complex entries appear all the time (they may even be the central objects in your instructor's dissertation) and have uses in physics and electrical engineering.

If  $A = [a_{ij}]$  is a matrix with complex entries, the *conjugate* is

$$\bar{A} = \begin{bmatrix} \overline{a_{11}} & \cdots & \overline{a_{1n}} \\ \vdots & & \vdots \\ \overline{a_{m1}} & \cdots & \overline{a_{mn}} \end{bmatrix}.$$

and the *conjugate transpose* (sometimes called the *adjoint* or *Hermitian transpose*) is obtained by first conjugating the matrix and then transposing it

$$\bar{A}^T = \begin{bmatrix} \overline{a_{11}} & \cdots & \overline{a_{m1}} \\ \vdots & & \vdots \\ \overline{a_{1n}} & \cdots & \overline{a_{nn}} \end{bmatrix}.$$

The conjugate-transpose is also sometimes denoted  $A^\dagger$  or  $A^*$ , depending on whether or not a physicist wrote the paper.

## 4.2 Determinants

**Definition.** The *determinant* (denoted  $\det$ ) is a function whose inputs are all of the columns of an  $n \times n$  real matrix  $A$ , whose output is a single real number, and which satisfies the following properties:

1.  $\det I_n = 1$ .
2.  $\det$  is a *multilinear* function (it is a linear transformation in each input).
3.  $\det$  is an *alternating* function (its sign changes whenever two columns are swapped).


*Remark.*

- The determinant is actually the unique such function with these properties.
- Some write  $|A|$  to mean  $\det A$ .


**How do we compute the determinant of a given matrix?**

### 4.2.1 Determinant of a $2 \times 2$ Matrix

The first property of the determinant is that  $\det(I_2) = 1$ , so we'd like to find a natural place where the value 1 naturally occurs relative to the columns of the identity matrix. Plotting the vectors  $\mathbf{e}_1$  and  $\mathbf{e}_2$ , we see that they naturally form a parallelogram (in particular, a square) of area 1, so maybe the natural choice is that the determinant should be the (signed) area of this parallelogram.



More generally, let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . For purposes of the picture, we will assume that  $a > c > 0$  and  $d > b > 0$ .



The (signed) area of the parallelogram is

$$\text{area} = (a+c)(b+d) - 2bc - cd - ab = ad - bc.$$

**Proposition 4.2.1.**  $\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc.$

*Proof.* By construction we know that this satisfies the property that  $\det I_2 = 1$ , so we only need to check the multilinearity and alternating conditions.

Multilinearity: it is a straightforward computation to show that

$$\det \begin{bmatrix} (a_1 + a_2) & b \\ (c_1 + c_2) & d \end{bmatrix} = \det \begin{bmatrix} a_1 & b \\ c_1 & d \end{bmatrix} + \det \begin{bmatrix} a_2 & b \\ c_2 & d \end{bmatrix}$$

and

$$\det \begin{bmatrix} a & (b_1 + b_2) \\ c & (d_1 + d_2) \end{bmatrix} = \det \begin{bmatrix} a & b_1 \\ c & d_1 \end{bmatrix} + \det \begin{bmatrix} a & b_2 \\ c & d_2 \end{bmatrix}$$

and that for any scalar  $k$ ,

$$\det \begin{bmatrix} ka & b \\ kc & d \end{bmatrix} = k \det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \det \begin{bmatrix} a & kb \\ c & kd \end{bmatrix}.$$

Alternating: it is a straightforward computation to show that

$$\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = -\det \begin{bmatrix} b & a \\ d & c \end{bmatrix}.$$

Since the determinant is the unique function with these properties, this must mean that our definition of determinant is correct.  $\square$

**Theorem 4.2.2.** For any  $n \times n$  matrix  $A$ ,  $\det A$  is the signed volume of the parallelepiped formed by the columns of  $A$ .

## 4.2.2 Determinant of a $n \times n$ Matrix

We already learned the determinant of a  $2 \times 2$  matrix. So how about for  $n \times n$  matrices where  $n > 2$ ? Computing the parallelepiped volume is a bit of a chore, but thankfully we can do it iteratively.

**Definition.** For an  $n \times n$  matrix  $A$ , the  $(i, j)$ -minor of  $A$ , denoted  $M_{i,j}$ , is the determinant of the *submatrix* formed by removing row  $i$  and column  $j$  from  $A$ .

**Example 4.2.3.** For  $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$ , the  $(2, 1)$ -minor is

$$M_{2,1} = \det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \det \begin{bmatrix} a_{12} & a_{13} \\ a_{32} & a_{33} \end{bmatrix} = a_{12}a_{33} - a_{32}a_{13}.$$

*Remark.* Your book uses the notation  $\det A_{ij}$  to denote the  $(i, j)$ -minor, but I think it just makes things more confusing.

**Theorem 4.2.4.** The determinant of an  $n \times n$  matrix  $A$  can be computed along the  $i^{\text{th}}$  row as the sum

$$\det A = \sum_{j=1}^n (-1)^{i+j} a_{ij} M_{i,j}.$$

or along the  $j^{\text{th}}$  column as the sum

$$\det A = \sum_{i=1}^n (-1)^{i+j} a_{ij} M_{i,j}.$$

It is an exercise to the reader to verify that the above sums satisfy the definition of the determinant and that they agree no matter along which row or column they are computed.

*Remark.* The quantity  $(-1)^{i+j} M_{i,j}$  is sometimes called the  $(i, j)$ -cofactor and the above sums are cofactor expansions.

**Example 4.2.5.** Compute  $\det A$  given  $A = \begin{bmatrix} 3 & 1 & 4 \\ -1 & 5 & -9 \\ 2 & 6 & -5 \end{bmatrix}$  along the first row.

$$\begin{aligned} \det A &= a_{11}M_{1,1} - a_{12}M_{1,2} + a_{13}M_{1,3} \\ &= 3 \det \begin{bmatrix} 3 & 1 & 4 \\ -1 & 5 & -9 \\ 2 & 6 & -5 \end{bmatrix} - 1 \det \begin{bmatrix} 3 & 1 & 4 \\ -1 & 5 & -9 \\ 2 & 6 & -5 \end{bmatrix} + 4 \det \begin{bmatrix} 3 & 1 & 4 \\ -1 & 5 & -9 \\ 2 & 6 & -5 \end{bmatrix} \\ &= 3 \det \begin{bmatrix} 5 & -9 \\ 6 & -5 \end{bmatrix} - \det \begin{bmatrix} -1 & -9 \\ 2 & -5 \end{bmatrix} + 4 \det \begin{bmatrix} -1 & 5 \\ 2 & 6 \end{bmatrix} \\ &= 3(29) - (23) + 4(-16) = 0. \end{aligned}$$

Having a zero determinant tells us that the corresponding parallelepiped has no volume, which means all three column vectors must live in the same plane (or even same line). So a determinant of zero is enough to tell us that the columns are linearly dependent!

**Proposition 4.2.6.** *For an  $n \times n$  matrix,  $\det(A) = 0$  if and only if the columns of  $A$  are linearly dependent.*

This means we can add onto the fundamental theorem of invertible matrices:

**Theorem 4.2.7.**  *$A$  is invertible if and only if  $\det(A) \neq 0$ .*

### 4.2.3 Properties of Determinants

**Definition.** A square matrix  $A = [a_{ij}]$  is *upper triangular* if  $a_{ij} = 0$  whenever  $i > j$ , and is *lower triangular* if  $a_{ij} = 0$  whenever  $i < j$ .

**Example 4.2.8.** Consider the following matrices

$$A = \begin{bmatrix} 1 & 2 & 3 \\ 0 & 4 & 5 \\ 0 & 0 & 6 \end{bmatrix}, \quad B = \begin{bmatrix} 1 & 0 & 0 \\ 2 & 3 & 0 \\ 4 & 5 & 6 \end{bmatrix}, \quad C = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}.$$

$A$  is upper triangular,  $B$  is lower triangular, and  $C$  (a diagonal matrix) is both upper and lower triangular.

**Theorem 4.2.9** (Poole Theorem 4.2). *If  $A$  is an  $n \times n$  triangular matrix, then  $\det(A) = a_{11}a_{22} \cdots a_{nn}$ , the product of the numbers along the diagonal.*

*Proof.* Perform the cofactor expansion along the first column (in the upper triangular case) or along the first row (in the lower triangular case).  $\square$

**Example 4.2.10.** Let's look at the matrices in Example 4.2.8. By computing the determinant of  $A$  along the first column, we have

$$\begin{aligned} \det A &= 1M_{1,1} - 0M_{2,1} + 0M_{3,1} \\ &= 1 \det \begin{bmatrix} 4 & 5 \\ 0 & 6 \end{bmatrix} = 1(4)(6). \end{aligned}$$

Now computing the determinant of  $B$  along the first row, we have

$$\begin{aligned} \det B &= 1M_{1,1} - 0M_{1,2} + 0M_{1,3} \\ &= 1 \det \begin{bmatrix} 3 & 0 \\ 5 & 6 \end{bmatrix} = 1(3)(6). \end{aligned}$$

Finally, computing the determinant of  $C$  along the first row (although the first column is perfectly fine as well),

$$\begin{aligned} \det C &= 1M_{1,1} - 0M_{1,2} + 0M_{1,3} \\ &= 1 \det \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix} = 1(2)(3). \end{aligned}$$

**Theorem 4.2.11** (Poole Theorems 4.7 - 4.10). *If  $A$  and  $B$  are  $n \times n$  matrices and  $k$  is a scalar, then*

1.  $\det(AB) = (\det A)(\det B)$ ,
2.  $\det(kA) = k^n(\det A)$ ,
3.  $\det A = \det A^T$ ,
4. *and if  $A$  is invertible,  $\det A^{-1} = \frac{1}{\det A}$ .*

**Example 4.2.12.** Verify each part of the theorem above with the following:

$$A = \begin{bmatrix} 3 & -1 \\ 8 & -2 \end{bmatrix}, \quad B = \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix}, \quad \text{and} \quad k = 5.$$

1.  $\det A = -6 + 8 = 2$ ,  $\det B = 4 - 1 = 3$ , and

$$\det(AB) = \det \begin{bmatrix} 5 & 1 \\ 14 & 4 \end{bmatrix} = 6 = (2)(3) = (\det A)(\det B).$$

2.  $\det kA = \det \begin{bmatrix} 15 & -5 \\ 40 & -10 \end{bmatrix} = 50 = 25(2) = k^2 \det A.$

3.  $\det A^T = \det \begin{bmatrix} 3 & 8 \\ -1 & -2 \end{bmatrix} = -6 + 8 = 2 = \det A$

4.  $\det A^{-1} = \det \begin{bmatrix} -1 & \frac{1}{2} \\ -4 & \frac{3}{2} \end{bmatrix} = -\frac{3}{2} + 2 = \frac{1}{2} = \frac{1}{\det A}$

### 4.3 Eigenvalues and Eigenvectors of $n \times n$ Matrices

We've already seen eigenvalues and eigenvectors for  $2 \times 2$  matrices, but now that we have defined determinants for  $n \times n$  matrices, we'll extend these definitions accordingly.

**Definition.** If  $A$  is a square matrix, then  $\det(A - \lambda I)$  is a polynomial with indeterminate  $\lambda$  and is called the *characteristic polynomial* of  $A$ . The *eigenvalues* of  $A$  are precisely the roots of the characteristic polynomial. For each eigenvalue  $\lambda$ , the corresponding *eigenspace* is  $E_\lambda = \text{Null}(A - \lambda I)$  and the nonzero vectors in  $E_\lambda$  are *eigenvectors*.

*Remark.* Non-square matrices do not have eigenvalues/eigenvectors, because if  $v \in \mathbb{R}^n$  is an eigenvector for  $A$ , then  $Av = \lambda v$  implies that  $v \in \mathbb{R}^n$  as well, hence  $A$  is  $n \times n$ . Non-square matrices have something called *singular values* which, in some sense, play the role of eigenvalues, but this is outside of the scope of this course.

**Example 4.3.1.** Find the eigenvalues and eigenvectors for  $A = \begin{bmatrix} 2 & 12 & 10 \\ 0 & -4 & -4 \\ 1 & 2 & 1 \end{bmatrix}$ .

We first compute  $\det(A - \lambda I)$  via cofactor expansion along the first column.

$$\begin{aligned} p(\lambda) = \det(A - \lambda I) &= \det \begin{bmatrix} 2 - \lambda & 12 & 10 \\ 0 & -4 - \lambda & -4 \\ 1 & 2 & 1 - \lambda \end{bmatrix} \\ &= (2 - \lambda) \det \begin{bmatrix} -4 - \lambda & -4 \\ 2 & 1 - \lambda \end{bmatrix} + 1 \det \begin{bmatrix} 12 & 10 \\ -4 - \lambda & -4 \end{bmatrix} \\ &= (2 - \lambda) ((-4 - \lambda)(1 - \lambda) + 8) + 1(-48 - 10(-4 - \lambda)) \\ &= -(\lambda^3 + \lambda^2 - 12\lambda) \\ &= -\lambda(\lambda + 4)(\lambda - 3) \end{aligned}$$

The characteristic polynomial factors nicely and the eigenvalues are  $-4, 0, 3$ . The corresponding eigenspaces are

$$\begin{aligned} E_{-4} &= \text{Null}(A + 4I) = \text{Null} \left( \begin{bmatrix} 6 & 12 & 10 \\ 0 & 0 & -4 \\ 1 & 2 & 5 \end{bmatrix} \right) = \text{Span} \left( \begin{bmatrix} -2 \\ 1 \\ 0 \end{bmatrix} \right), \\ E_0 &= \text{Null}(A - 0I) = \text{Null} \left( \begin{bmatrix} 2 & 12 & 10 \\ 0 & -4 & -4 \\ 1 & 2 & 1 \end{bmatrix} \right) = \text{Span} \left( \begin{bmatrix} 1 \\ -1 \\ 1 \end{bmatrix} \right), \\ E_3 &= \text{Null}(A - 3I) = \text{Null} \left( \begin{bmatrix} -1 & 12 & 10 \\ 0 & -7 & -4 \\ 1 & 2 & -2 \end{bmatrix} \right) = \text{Span} \left( \begin{bmatrix} 22 \\ -4 \\ 7 \end{bmatrix} \right). \end{aligned}$$

**Example 4.3.2.** Consider the matrix  $A = \begin{bmatrix} 3 & -2 & -4 \\ 0 & 1 & 2 \\ 0 & 0 & 1 \end{bmatrix}$ . The characteristic polynomial for  $A$  is

$$p(\lambda) = (3 - \lambda)(1 - \lambda)^2$$



hence the eigenvalues are 1, 3. The corresponding eigenspaces are

$$E_3 = \text{Span} \left( \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right) \quad \text{and} \quad E_1 = \text{Span} \left( \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix} \right).$$

Notice that the above  $3 \times 3$  matrix had only two distinct eigenvalues, but if we counted multiplicity as well ( $(1 - \lambda)$  appears with multiplicity 2 in the characteristic polynomial), then in fact we have exactly 3 eigenvalues. It may also be interesting to notice that  $\dim(E_1) = \dim(E_3)$ .

**Definition.** The *algebraic multiplicity* of an eigenvalue  $\lambda$  is the multiplicity as a root of the characteristic polynomial, and the *geometric multiplicity* is the dimension of the eigenspace  $E_\lambda$ .

These two different notions of multiplicity will be important in the next section. We'll note that if  $A$  is an  $n \times n$  matrix, then the sum of all of the algebraic multiplicities will always be  $n$ .

**Theorem 4.3.3** (Poole - Theorem 4.15). *If  $A$  is a triangular (or diagonal) matrix, then the eigenvalues are precisely the entries appearing along the diagonal.*

*Proof.* If  $A$  is triangular, then by Theorem 4.2.9

$$p(\lambda) = \det(A - \lambda I) = (a_{11} - \lambda)(a_{22} - \lambda) \cdots (a_{nn} - \lambda).$$

□

**Theorem 4.3.4.** *If  $A$  is an  $n \times n$  square matrix with eigenvalues  $\lambda_1, \dots, \lambda_n$  (not necessarily all distinct), then  $\det A = \lambda_1 \lambda_2 \cdots \lambda_n$ , the product of all of the eigenvalues.*

*Proof.* The above theorem is easy to see in the case that  $A$  is triangular - the product of the roots of a polynomial are precisely the constant term, and the constant term of the characteristic polynomial is exactly  $a_{11} a_{22} \cdots a_{nn}$ . □

From this it follows that we have yet another test for invertibility:

**Theorem 4.3.5.** *A square matrix  $A$  is invertible if and only if 0 is not an eigenvalue of  $A$ .*

### 4.3.1 Relationship to Matrix Operations

It is natural to ask about the interplay between eigenvalues/eigenvectors and matrix operations like inversion and exponentiation.

**Example 4.3.6.** Consider  $A = \begin{bmatrix} 1 & -2 \\ 1 & 4 \end{bmatrix}$ . The characteristic polynomial is

$$p_A(\lambda) = (2 - \lambda)(3 - \lambda)$$

hence the eigenvalues are 2, 3 and the corresponding eigenspaces are

$$E_2 = \text{Span} \left( \begin{bmatrix} -2 \\ 1 \end{bmatrix} \right) \quad \text{and} \quad E_3 = \text{Span} \left( \begin{bmatrix} -1 \\ 1 \end{bmatrix} \right).$$

With  $A$  as above, we have that  $A^{-1} = \frac{1}{6} \begin{bmatrix} 4 & 2 \\ -1 & 1 \end{bmatrix}$  and the eigenvalues are  $\frac{1}{2}, \frac{1}{3}$  – reciprocals of  $A$ 's eigenvalues. What's more, notice that

$$A^{-1} \begin{bmatrix} -2 \\ 1 \end{bmatrix} = \frac{1}{2} \begin{bmatrix} -2 \\ 1 \end{bmatrix} \quad \text{and} \quad A^{-1} \begin{bmatrix} -1 \\ 1 \end{bmatrix} = \frac{1}{3} \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

so the reciprocal eigenvalues of  $A^{-1}$  have the same eigenvectors as the eigenvalues of  $A$ ! With  $A$  as before, we have that  $A^2 = \begin{bmatrix} -1 & -10 \\ 5 & 14 \end{bmatrix}$  and the eigenvalues are 4, 9 – squares of  $A$ 's eigenvalues. What's more, notice that


$$A^2 \begin{bmatrix} -2 \\ 1 \end{bmatrix} = 2^2 \begin{bmatrix} -2 \\ 1 \end{bmatrix} \quad \text{and} \quad A^2 \begin{bmatrix} -1 \\ 1 \end{bmatrix} = 3^2 \begin{bmatrix} -1 \\ 1 \end{bmatrix}$$

so the squared eigenvalues of  $A^2$  have the same eigenvectors as the eigenvalues of  $A$ !


**Theorem 4.3.7.** *Let  $A$  be a square matrix with eigenvalue  $\lambda$  and corresponding eigenvector  $\mathbf{v}$ .*

1. *For any positive integer  $n$ ,  $\lambda^n$  is an eigenvalue of  $A^n$  with corresponding eigenvector  $\mathbf{v}$ .*
2. *If  $A$  is invertible, then  $\frac{1}{\lambda}$  is an eigenvalue of  $A^{-1}$  with corresponding eigenvector  $\mathbf{v}$ .*

The above theorem also makes sense geometrically. Each application of the transformation  $A$  stretches its eigenvector  $\mathbf{v}$  by a factor of  $\lambda$ :



Similarly, each application of  $A^{-1}$  “undoes” the stretching of its eigenvector  $\mathbf{v}$  by a factor of  $\lambda$  (i.e., stretches instead by a factor of  $\frac{1}{\lambda}$ ): The above theorem also makes sense geometrically. Each application of  $A$  stretches its eigenvector by a factor of  $\lambda$



## 4.4 Similarity and Diagonalization

As we saw in the previous section, triangular and diagonal matrices were very nice from a computational standpoint, so it would be nice to convert a matrix into triangular form in a meaningful way. We already know that we can do this with row reduction, but this process does not preserve eigenvalues (any invertible matrix row reduces to the identity, for example), so in this section we will look at another process that does retain the useful eigen-information.

**Definition.** Two  $n \times n$  matrices  $A$  and  $B$  are called *similar* if there is an invertible  $n \times n$  matrix  $P$  for which  $P^{-1}AP = B$ . We sometimes write “ $A \sim B$ ” to mean “ $A$  is similar to  $B$ .” We also sometimes refer to the product  $P^{-1}AP$  as “conjugation of  $A$  by  $P$ .”

*Remark.* Such a  $P$  is not unique. For example,  $P^{-1}IP = I$  is true for every invertible matrix  $P$ .

**Example 4.4.1.**  $A = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 0 \\ 0 & 5 \end{bmatrix}$  are similar. With  $P = \begin{bmatrix} -3 & 1 \\ 1 & 1 \end{bmatrix}$ , we have

$$\begin{aligned} P^{-1}AP &= \begin{bmatrix} -3 & 1 \\ 1 & 1 \end{bmatrix}^{-1} \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} -3 & 1 \\ 1 & 1 \end{bmatrix} \\ &= \frac{1}{-4} \begin{bmatrix} 1 & -1 \\ -1 & -3 \end{bmatrix} \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix} \begin{bmatrix} -3 & 1 \\ 1 & 1 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & 5 \end{bmatrix} = B \end{aligned}$$

**Example 4.4.2.** The matrices  $A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  are not similar.

If they were, we could find a matrix  $P = \begin{bmatrix} x & y \\ z & w \end{bmatrix}$  for which  $B = P^{-1}AP$ . In this case, we would have

$$\begin{aligned} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} &= \frac{1}{xw - yz} \begin{bmatrix} w & -y \\ -z & x \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} x & y \\ z & w \end{bmatrix} \\ &= \frac{1}{xw - yz} \begin{bmatrix} wx + yz & 2wy \\ -2xz & -wx - yz \end{bmatrix} \end{aligned}$$

and it's impossible that both  $(wx + yz) = 1$  and  $(-wx - yz) = 1$  (otherwise  $wx + yz = -(wx + yz)$ , hence  $wx + yz = 0$ ).

### 4.4.1 Properties of similarity and similar matrices

**Theorem 4.4.3** (Poole Theorem 4.21). *Let  $A, B, C$  be  $n \times n$  matrices.*

- reflexive:  $A \sim A$ .
- symmetric: *If  $A \sim B$  then  $B \sim A$ .*
- transitive: *If  $A \sim B$  and  $B \sim C$  then  $A \sim C$ .*

Each of the following properties are easily verified, say with the matrices from Example 4.4.1.

**Theorem 4.4.4** (Poole Theorem 4.22). *Let  $A$  and  $B$  be similar  $n \times n$  matrices. Then*

- a.  $\det A = \det B$
- b.  $A$  is invertible if and only if  $B$  is invertible.
- c.  $A$  and  $B$  have the same rank.
- d.  $A$  and  $B$  have the same characteristic polynomial.
- e.  $A$  and  $B$  have the same eigenvalues.
- f.  $A^m \sim B^m$  for any positive integer  $m$ .

*Partial proof sketch.* Let  $P$  be a matrix for which  $B = P^{-1}AP$ .

- a.  $\det B = \det(P^{-1}AP) = (\det P^{-1})(\det A)(\det P) = \left(\frac{1}{\det P}\right)(\det A)(\det P) = \det A$
- b.  $B^{-1} = (P^{-1}AP)^{-1} = P^{-1}A^{-1}(P^{-1})^{-1} = P^{-1}A^{-1}P$
- c. This follows from the fact that for any invertible matrix  $P$ ,  $\text{Rank}(A) = \text{Rank}(PA) = \text{Rank}(AP)$ .
- d.  $\det(B - \lambda I) = \det(P^{-1}AP - \lambda P^{-1}IP) = \det(P^{-1}(A - \lambda I)P) = \det(A - \lambda I)$
- e.  $A$  and  $B$  have the same characteristic polynomials
- f.  $B^m = \underbrace{(P^{-1}AP) \cdots (P^{-1}AP)}_m = P^{-1} \underbrace{A \cdots A}_m P = P^{-1}A^mP$

□

To check whether two given matrices  $A$  and  $B$  are similar requires finding the matrix  $P$  satisfying  $P^{-1}AP = B$ , which as we saw from 4.4.2, could be quite laborious. The above theorem is actually most useful for showing that two matrices are not similar (in fact, no single part of the theorem is enough to deduce that two matrices are similar).

**Example 4.4.5.** The matrices  $A$  and  $B$  in Example 4.4.2 are not similar because  $\det A = -1$  and  $\det B = 1$ .

**Example 4.4.6.** Although matrices  $A = \begin{bmatrix} 2 & 0 \\ 0 & 3 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 0 \\ 0 & 6 \end{bmatrix}$  have the same rank ( $\text{Rank } A = \text{Rank } B = 2$ ) and determinant ( $\det A = \det B = 6$ ), they are not similar because their eigenvalues and characteristic polynomials are different.

## 4.4.2 Diagonalization

**Definition.** A matrix  $A$  is *diagonalizable* if it is similar to a diagonal matrix  $D$ , i.e. if there is some invertible matrix  $P$  so that  $P^{-1}AP = D$ .

**Example 4.4.7.** From Example 4.4.1,  $A = \begin{bmatrix} 2 & 3 \\ 1 & 4 \end{bmatrix}$  is diagonalizable since it is similar to the diagonal matrix  $B = \begin{bmatrix} 1 & 0 \\ 0 & 5 \end{bmatrix}$ .

Notice that the characteristic polynomial of  $A$  above is

$$\det(A - \lambda I) = (2 - \lambda)(4 - \lambda) - 3 = \lambda^2 - 6\lambda + 5 = (\lambda - 1)(\lambda - 5)$$

and thus  $B$  contains  $A$ 's eigenvalues along the diagonal. This gives us a clue as to how one can go about finding the matrix  $P$  used to conjugate  $A$  into a diagonal matrix (if possible).

**Theorem 4.4.8** (Poole Theorem 4.23). *Let  $A$  be an  $n \times n$  matrix. Then  $A$  is diagonalizable if and only if  $A$  has  $n$  linearly independent eigenvectors.*

*More precisely,  $D = P^{-1}AP$  if and only if the columns of  $P$  are the eigenvectors of  $A$  and if the  $(i, i)$  entry of  $D$  is the eigenvalue corresponding to the  $i^{\text{th}}$  column of  $P$ .*

I won't sketch the proof, but the core observation is the following:

If  $P^{-1}AP = D$ , then this rearranges to  $AP = PD$ . So if  $\mathbf{p}_i$  is the  $i^{\text{th}}$  column of  $P$  and  $\lambda_i$  is the  $(i, i)$  entry in  $D$ , then

$$\begin{aligned}
 AP &= PD \\
 A \begin{bmatrix} \mathbf{p}_1 & \cdots & \mathbf{p}_n \end{bmatrix} &= \begin{bmatrix} \mathbf{p}_1 & \cdots & \mathbf{p}_n \end{bmatrix} \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix} \\
 \begin{bmatrix} A\mathbf{p}_1 & & A\mathbf{p}_n \end{bmatrix} &= \begin{bmatrix} \lambda_1\mathbf{p}_1 & & \lambda_n\mathbf{p}_n \end{bmatrix}
 \end{aligned}$$

and so  $A\mathbf{p}_i = \lambda_i\mathbf{p}_i$ , hence the  $\lambda_i$  are eigenvalues for  $A$  with corresponding eigenvectors  $\mathbf{p}_i$ .

**Example 4.4.9.** Let  $A = \begin{bmatrix} 2 & 3 \\ 0 & 2 \end{bmatrix}$ . The characteristic polynomial for  $A$  is

$$\det(A - \lambda I) = \det \begin{bmatrix} 2 - \lambda & 3 \\ 0 & 2 - \lambda \end{bmatrix} = (2 - \lambda)^2$$

so  $A$  has a single eigenvalue of 2 with algebraic multiplicity 2. The corresponding eigenspace is

$$E_2 = \text{Null}(A - 2I) = \text{Span} \left( \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right).$$

and so the eigenvalue 2 has geometric multiplicity 1. This means that there are not enough linearly independent eigenvectors to form our invertible matrix  $P$  (the one for which  $P^{-1}AP$  is a diagonal matrix), hence  $A$  is not diagonalizable.

**Example 4.4.10.** Consider the  $3 \times 3$  matrix  $A = \begin{bmatrix} 4 & -3 & -3 \\ 3 & -2 & -3 \\ -1 & 1 & 2 \end{bmatrix}$ . The characteristic polynomial is

$$\det(A - \lambda I) = \det \begin{bmatrix} 4 - \lambda & -3 & -3 \\ 3 & -2 - \lambda & -3 \\ -1 & 1 & 2 - \lambda \end{bmatrix} = -(\lambda - 1)^2(\lambda - 2)$$

and the eigenvalues are 1 and 2 (with algebraic multiplicities 2 and 1, respectively). The corresponding eigenspaces are

$$E_1 = \text{Null}(A - I) = \text{Span} \left( \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right) \quad E_2 = \text{Null}(A - 2I) = \text{Span} \left( \begin{bmatrix} -3 \\ -3 \\ 1 \end{bmatrix} \right)$$

and so the eigenvalues 1 and 2 have geometric multiplicities 2 and 1 (respectively). It is readily seen that the vectors we used to define  $E_1$  are linearly independent, so the following matrix is invertible:

$$P = \begin{bmatrix} 1 & 1 & -3 \\ 1 & 0 & -3 \\ 0 & 1 & 1 \end{bmatrix}.$$

We then diagonalize  $A$ :

$$P^{-1}AP = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 2 \end{bmatrix}.$$

What these examples have highlighted is that a matrix may only fail to be diagonalizable if it has repeated eigenvalues.

**Theorem 4.4.11** (Poole Theorem 4.24). *Let  $A$  be an  $n \times n$  matrix with distinct eigenvalues  $\lambda_1, \dots, \lambda_k$ . Let  $\mathcal{B}_i$  be the basis for  $E_{\lambda_i}$ . The union of the  $\mathcal{B}_i$ 's (i.e. the collection of all basis vectors in the  $\mathcal{B}_i$ 's) is a linearly independent set.*

**Corollary 4.4.12** (Poole Theorem 4.25). *If  $A$  is an  $n \times n$  matrix with  $n$  distinct eigenvalues, then  $A$  is diagonalizable.*

What is it about the repeated eigenvalues that causes the failure of diagonalizability of a matrix  $A \in \mathbb{R}^{n \times n}$ ? Well, we need there to be  $n$  linearly independent eigenvectors, so we need the geometric multiplicity for each eigenvalue to be as large as possible.

**Lemma 4.4.13** (Poole Lemma 4.26). *The geometric multiplicity of an eigenvalue  $\lambda$  is less than or equal to its algebraic multiplicity.*

All of this culminates in the following result:

**Theorem 4.4.14** (Diagonalization Theorem). *Let  $A$  be an  $n \times n$  matrix with distinct eigenvalues  $\lambda_1, \dots, \lambda_k$ . The following are equivalent:*

a.  $A$  is diagonalizable.

b. The union of the basis vectors from each  $E_{\lambda_i}$  is a set of  $n$  vectors. In other words,  $n = \sum_{i=1}^k \dim(E_{\lambda_i})$ .

c. For each  $i$ , the algebraic multiplicity of  $\lambda_i$  is equal to the geometric multiplicity of  $\lambda_i$ .

**Example 4.4.15.** The matrix  $A = \begin{bmatrix} 3 & -1 & 2 \\ 3 & -1 & 6 \\ -2 & 2 & -2 \end{bmatrix}$  has characteristic polynomial

$$\det(A - \lambda I) = (2 - \lambda)^2(4 - \lambda).$$

The eigenvalue 2 has geometric multiplicity 2 and the eigenvalue 4 has geometric multiplicity 1. By the Diagonalization Theorem,  $A$  is diagonalizable – you can verify that an appropriate conjugating matrix is

$$P = \begin{bmatrix} -1 & -2 & 1 \\ -3 & 0 & 1 \\ 2 & 1 & 0 \end{bmatrix}.$$

**Example 4.4.16.** The matrix  $A = \begin{bmatrix} 1 & 0 & 2 \\ 0 & 3 & 4 \\ 0 & 0 & 3 \end{bmatrix}$  has characteristic polynomial

$$\det(A - \lambda I) = (1 - \lambda)(3 - \lambda)^2.$$

Both eigenvalues 1 and 3 have geometric multiplicity 1, so by the Diagonalization Theorem,  $A$  is not diagonalizable.

### 4.4.3 Computational power of diagonal matrices

Notice that for a diagonal matrix  $D = \text{diag}(d_1, \dots, d_n)$  and any positive integer  $k$ ,

$$D^k = \begin{bmatrix} d_1^k & & \\ & \ddots & \\ & & d_n^k \end{bmatrix}.$$

Moreover, if  $D$  is invertible, then

$$D^{-k} = \begin{bmatrix} d_1^{-k} & & \\ & \ddots & \\ & & d_n^{-k} \end{bmatrix} = \begin{bmatrix} \frac{1}{d_1^k} & & \\ & \ddots & \\ & & \frac{1}{d_n^k} \end{bmatrix}.$$

This is instantaneous. For a general  $n \times n$  matrix  $A$ , computing  $A^k$  in the usual way is *extremely* computationally expensive. However, if  $A$  is diagonalizable, we can write  $P^{-1}AP = D$ , hence

$$D^k = (P^{-1}AP)^k = P^{-1}A^kP \quad \implies \quad A^k = PD^kP^{-1}.$$

In this way, computing the  $k^{\text{th}}$  power of  $A$  is only as computationally difficult as diagonalizing  $A$ .

**Example 4.4.17.** Let  $A = \begin{bmatrix} 11 & -6 \\ 15 & -8 \end{bmatrix}$ . One can readily check that  $A$  has eigenvalues 1, 2, hence is diagonalizable (and since the eigenvalues are all nonzero,  $A$  is invertible). Through the usual methods, we can obtain

$$A = \begin{bmatrix} 3 & 2 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 5 & 3 \end{bmatrix}^{-1}$$

whence, for any integer  $k$ ,

$$A^k = \begin{bmatrix} 3 & 2 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 1^k & 0 \\ 0 & 2^k \end{bmatrix} \begin{bmatrix} 3 & 2 \\ 5 & 3 \end{bmatrix}^{-1} = \begin{bmatrix} 3 & 2 \\ 5 & 3 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & 2^k \end{bmatrix} \begin{bmatrix} -3 & 2 \\ 5 & -3 \end{bmatrix} = \begin{bmatrix} -9 + 10(2^k) & 6 - 6(2^k) \\ -15 + 15(2^k) & 10 - 9(2^k) \end{bmatrix}.$$

Similarly,

$$A^{-2} = \begin{bmatrix} -9 + 10(2^{-2}) & 6 - 6(2^{-2}) \\ -15 + 15(2^{-2}) & 10 - 9(2^{-2}) \end{bmatrix} = \begin{bmatrix} -13/2 & 9/2 \\ -45/4 & 31/4 \end{bmatrix}$$



## 1.2 Length and Angle: The Dot Product

Recall the following

**Definition.** For vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ , the *dot product* of  $\mathbf{u}$  and  $\mathbf{v}$ , denoted  $\mathbf{u} \cdot \mathbf{v}$  is

$$\mathbf{u} \cdot \mathbf{v} = u_1v_1 + \cdots + u_nv_n.$$

*Remark.* When  $\mathbf{u}, \mathbf{v}$  are written as column vectors, the product

$$\mathbf{u}^T \mathbf{v} = [u_1v_1 + \cdots + u_nv_n]$$

is a vector in  $\mathbb{R}^1$ , so by identifying  $\mathbb{R}^1$  with  $\mathbb{R}$ , we have that  $\mathbf{u}^T \mathbf{v} = \mathbf{u} \cdot \mathbf{v}$  (the product  $\mathbf{u}^T \mathbf{v}$  is called an *inner product*).

Recall also the following facts about the dot product

**Theorem 1.2.1** (Poole Theorem 1.2). *Let  $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^n$  and let  $k$  be some scalar. Then*


1.  $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$
2.  $\mathbf{u} \cdot (\mathbf{v} + \mathbf{w}) = (\mathbf{u} \cdot \mathbf{v}) + (\mathbf{u} \cdot \mathbf{w})$
3.  $(\mathbf{v} + \mathbf{w}) \cdot \mathbf{u} = (\mathbf{v} \cdot \mathbf{u}) + (\mathbf{w} \cdot \mathbf{u})$
4.  $(k\mathbf{u}) \cdot \mathbf{v} = \mathbf{u} \cdot (k\mathbf{v}) = k(\mathbf{u} \cdot \mathbf{v})$
5. For every  $\mathbf{u}$  we have that  $\mathbf{u} \cdot \mathbf{u} \geq 0$ , with equality if and only if  $\mathbf{u} = \mathbf{0}$ .

### 1.2.1 Length

Notice that for a vector  $\mathbf{v} = [x, y] \in \mathbb{R}^2$ ,

$$\mathbf{v} \cdot \mathbf{v} = x^2 + y^2,$$

which, from the Pythagorean theorem, is precisely the square of the length of  $\mathbf{v}$ .



**Definition.** The *length* or *norm* of a vector  $\mathbf{v} \in \mathbb{R}^n$  is the scalar defined by

$$\|\mathbf{v}\| = \sqrt{\mathbf{v} \cdot \mathbf{v}} = \sqrt{v_1^2 + v_2^2 + \cdots + v_n^2}$$

The following are immediate consequences of the properties of the dot product in Theorem 1.2.1


**Theorem 1.2.2** (Poole Theorem 1.3). For  $\mathbf{v} \in \mathbb{R}^n$  and a scalar  $k$ ,

1.  $\|\mathbf{v}\| = 0$  if and only if  $\mathbf{v} = \mathbf{0}$ .
2.  $\|k\mathbf{v}\| = |k|\|\mathbf{v}\|$ .

The following follows from the classical geometry result of the same name.

**Theorem 1.2.3** (Triangle Inequality). For  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ ,

$$\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|.$$




**Definition.** A vector  $\mathbf{v}$  is called a *unit vector* if  $\|\mathbf{v}\| = 1$ .

*Remark.* Every unit vector in  $\mathbb{R}^2$  corresponds to a point on the unit circle. Every unit vector in  $\mathbb{R}^3$  corresponds to a point on the unit sphere. Generally, every unit vector in  $\mathbb{R}^n$  corresponds to a point on the unit  $(n - 1)$ -sphere.

Let  $\mathbf{v}$  be any nonzero vector and let  $\ell = \|\mathbf{v}\|$  be its length. Then the vector  $\frac{\mathbf{v}}{\ell}$  is a unit vector because

$$\left\| \frac{\mathbf{v}}{\ell} \right\| = \frac{\|\mathbf{v}\|}{\ell} = \frac{\ell}{\ell} = 1$$

The process above is called *normalization*, and it always produces a vector in the same direction as  $\mathbf{v}$  but with unit length.



*Remark.* If  $\|\mathbf{v}\| > 1$ , then normalization corresponds to shrinking  $\mathbf{v}$  (pictured above), but if  $\|\mathbf{v}\| < 1$ , then normalization stretches  $\mathbf{v}$ .

*Remark.* Despite the similarities in name, “normalization” is unrelated to the concept of a “normal vector.” What you’ll find is that “normal” is probably the most over-used word in mathematics. Because there aren’t any around me as I type this, I’m going to go ahead and blame the physicists for the abuse of language.

### 1.2.2 Distances

Recall that, for two points  $P(x_1, y_1)$  and  $Q(x_2, y_2)$  in the plane, we have that the distance between them is given by


$$d(P, Q) = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$

If we identify the point  $P(x_1, y_1)$  with the vector  $\mathbf{u} = [x_1, y_1]$  and the point  $Q(x_2, y_2)$  with the vector  $\mathbf{v} = [x_2, y_2]$ , then the right-hand side of the equation is just  $\|\mathbf{u} - \mathbf{v}\|$ . As such, we can define distances between vectors using the obvious analog.

**Definition.** Given two vectors  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ , the *distance* between  $\mathbf{u}$  and  $\mathbf{v}$  is


$$d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\|.$$

*Remark.* Visualizing vectors as arrows emanating from the origin, distance, as above, is actually measuring the distance between the heads of the arrows.



### 1.2.3 Projections

Up until now we haven't really given any geometric description of what the dot product actually measures, so let's consider  $\mathbf{e}_1 = [1, 0]$  and  $\mathbf{v} = [x, y]$ . A straightforward computation shows that  $\mathbf{e}_1 \cdot \mathbf{v} = x$ . Looking at the picture below



We see that  $x$  is precisely the length of the “shadow” cast by  $\mathbf{v}$  on the line spanned by  $\mathbf{e}_1$ . More generally, for two vectors  $\mathbf{u}$  and  $\mathbf{v}$ , the dot product  $\mathbf{u} \cdot \mathbf{v}$  is the length of “shadow” of  $\mathbf{v}$  cast onto  $\mathbf{u}$ , but then scaled by the length of  $\mathbf{v}$  (by symmetry of the dot product, this description also works *mutatis mutandis* for the “shadow” cast by  $\mathbf{u}$  onto  $\mathbf{v}$ ).

Let's formalize this notion of a “shadow” cast by  $\mathbf{v}$  onto  $\mathbf{u}$  keeping in mind two things:


1.  $\mathbf{u}$  should be normalized to be a unit vector so as to preserve the length of  $\mathbf{v}$ 's shadow, and
2. the shadow of a vector should also be a vector in the direction of  $\mathbf{u}$  (i.e. a scalar multiple of  $\mathbf{u}$ ).

**Definition.** Given two vectors  $\mathbf{u}$  and  $\mathbf{v}$ , the *projection* (or *orthogonal projection*) of  $\mathbf{v}$  onto  $\mathbf{u}$  is the vector  $\text{proj}_{\mathbf{u}}(\mathbf{v})$  given by

$$\text{proj}_{\mathbf{u}}(\mathbf{v}) = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\|^2} \right) \mathbf{u} = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\mathbf{u} \cdot \mathbf{u}} \right) \mathbf{u}.$$

**Example 1.2.4.** Let  $\mathbf{u} = [4, -1]$  and  $\mathbf{v} = [3, 5]$ . Using the formula from the definition we have

$$\text{proj}_{\mathbf{u}}(\mathbf{v}) = \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\mathbf{u} \cdot \mathbf{u}} \right) \mathbf{u} = \left( \frac{7}{17} \right) \mathbf{u} = \left[ \frac{28}{17}, \frac{-7}{17} \right]$$



Notice that the vector  $\mathbf{v} - \text{proj}_{\mathbf{u}}(\mathbf{v})$  (represented by a dashed line) is perpendicular to  $\mathbf{u}$ . Perpendicular vectors are called *orthogonal* (hence the name “orthogonal projection”). What’s more

$$\begin{aligned} (\mathbf{v} - \text{proj}_{\mathbf{u}}(\mathbf{v})) \cdot \mathbf{u} &= \mathbf{v} \cdot \mathbf{u} - \left( \frac{\mathbf{u} \cdot \mathbf{v}}{\mathbf{u} \cdot \mathbf{u}} \right) \mathbf{u} \cdot \mathbf{u} \\ &= \mathbf{v} \cdot \mathbf{u} - \mathbf{u} \cdot \mathbf{v} \\ &= \mathbf{u} \cdot \mathbf{v} - \mathbf{u} \cdot \mathbf{v} \\ &= 0. \end{aligned}$$

So we deduce from this the following fact

**Theorem 1.2.5.** *Two vectors  $\mathbf{u}, \mathbf{v}$  in  $\mathbb{R}^n$  are orthogonal if and only if  $\mathbf{u} \cdot \mathbf{v} = 0$ .*

## 5.1 Orthogonality in $\mathbb{R}^n$

## 5.2 Orthogonal Complements and Orthogonal Projections

### 5.3 The Gram-Schmidt Process and the $QR$ Factorization



## 7.3 Least Squares